



Chemical and ruminal in vitro evaluation of Canadian canola meals produced over 4 years¹

Glen A. Broderick,^{*2,3} Stefania Colombini,[†] Sara Costa,[†] Mehmet A. Karsli,[‡] and Antonio P. Faciola[§]

^{*}Agricultural Research Service, USDA, US Dairy Forage Research Center, 1925 Linden Drive, Madison, WI 53706

[†]Dipartimento di Scienze Animali, Università degli Studi di Milano, Milan, 20133 Italy

[‡]Department of Animal Nutrition, Veterinary Faculty, Kirikkale University, 71451 Turkey

[§]Department of Agriculture, Nutrition, and Veterinary Science, University of Nevada, Reno 89557

ABSTRACT

To test the effects of year and processing plant on the nutritional value of canola meal (CM), 3 CM samples/yr were collected from each of 12 Canadian production plants over 4 yr (total = 144). Samples of CM were analyzed for differences in chemical composition and for in vitro ruminal protein degradability using the Michaelis-Menten inhibitor in vitro (MMIIV) method. In the MMIIV method, protein degradation rate (k_d) was estimated by 2 methods: from net release (i.e., blank corrected) of (1) ammonia plus AA determined by *o*-phthaldialdehyde fluorescence (OPA_F) assay or (2) ammonia, AA, plus oligopeptides determined by *o*-phthaldialdehyde absorbance (OPA_A) assay; rumen-undegradable protein (RUP) was computed assuming passage rates of 0.16 and 0.06/h for, respectively, soluble and insoluble protein. Casein, solvent soybean meal (SSBM), and expeller soybean meal (ESBM) were included in all incubations as standard proteins. Differences among years and plants were assessed using the mixed procedures of SAS. Small but significant differences were found in CM among years for chemical composition, including N solubility; some of these differences may have been related to changes in our analytical methods over time. However, adjustment of degradation activity of individual in vitro incubations based on the mean degradation activity over all incubations yielded k_d and RUP that did not differ by year using either assay. Simultaneously incubating CM samples from 2 yr in the same in vitro runs confirmed that no year effects existed for k_d or RUP. Differences

existed in chemical composition of CM among the 12 processing plants over the 4 yr of sample collection. Moreover, consistent differences in k_d and RUP were observed among plants: k_d ranged from 0.069 to 0.113/h (OPA_A assay) and 0.075 to 0.120/h (OPA_F assay), and RUP estimates ranged from 51 to 43% (OPA_A assay) and 49 to 41% (OPA_F assay). Regression of k_d on insoluble N content of CM yielded correlation coefficients (R^2) = 0.40 (OPA_A assay) and 0.42 (OPA_F assay), and regressions of k_d on NDIN and N-fraction B₃ yielded $R^2 < 0.02$. Mean estimates from both OPA_A and OPA_F assays for casein, SSBM, ESBM, and CM were, respectively, k_d = 0.764, 0.161, 0.050, and 0.093/h and RUP = 18, 33, 56, and 45%. A range of 8 percentage units from lowest to highest RUP suggests that substantial differences exist in metabolizable protein content of CM produced by different processing plants.

Key words: canola meal, chemical composition, ruminal degradation, rumen-undegraded protein

INTRODUCTION

Increased production of canola has resulted in greater availability of canola meal (CM) as an alternative to soybean meal (SBM) for protein supplementation of lactating dairy cows (Hickling, 2008). Meta-analyses of published findings showed that replacing SBM with CM significantly increased milk protein yield (Martineau et al., 2013) and increased feed intake and yield of milk and milk components (Huhtanen et al., 2011). We observed numeric increases in milk and protein yield when CM replaced supplemental protein from SBM in 16.5% CP diets in dairy cows (Brito and Broderick, 2007). Brito et al. (2007) found that the proportion of RUP in CM was numerically greater than that in SBM. Huhtanen et al. (2011) also concluded that CM contributed amounts of RUP and MP that were at least equal to SBM. Ruminal in situ studies conducted by Maxin et al. (2013a) showed that SBM had a more rapid degradation rate, higher effective degradability, and lower RUP than CM. More recently (Broderick et al., 2015),

Received February 8, 2016.

Accepted June 6, 2016.

¹Mention of any trademark or proprietary product in this paper does not constitute a guarantee or warranty of the product by the USDA or the Agricultural Research Service and does not imply its approval to the exclusion of other products that also may be suitable.

²Retired; present address: Broderick Nutrition and Research LLC, 221 Glen Hollow Road, Madison, WI 53705.

³Corresponding author: gbroderi@wisc.edu

increased DMI and yield of milk and milk protein were associated with reduced ruminal concentrations of ammonia and branched-chain VFA in cows fed CM versus SBM, suggesting lower ruminal degradation of CM protein. The National Research Council (NRC, 2001) model indicates ruminal protein degradation rates of 7.5%/h for 48% SBM and 10.4%/h for CM, and RUP values of 43% for 48% SBM and 36% for CM (at DMI = 4% of BW with 50% dietary DM fed as forage). These NRC (2001) data appear to be inconsistent with the greater RUP in CM reported by Maxin et al. (2013a) and the reduced ruminal ammonia and branched-chain VFA concentrations observed by Broderick et al. (2015) when CP from CM replaced equal CP from SBM.

Other evidence indicates that CM may be a more effective protein supplement than certain byproduct feeds such as distillers dried grains plus solubles (**DDGS**): Swanepoel et al. (2014) observed that replacing corn DDGS with CM increased both milk and true protein yield. Although milk and protein yield were not different, Acharya et al. (2015) found that replacing corn DDGS with CM significantly improved efficiency of MP utilization. Mutsvangwa et al. (2016) reported that substituting CM for wheat DDGS gave a numeric yield increase of 1.1 kg of milk/d plus increased omasal flow of Thr and Trp and tended to increase omasal flow of His and Lys.

Growing conditions experienced in canola production vary substantially from year to year, and we speculated that these differences might alter the nutritional quality of CM for ruminants. Therefore, the objectives of this study were to (1) determine if year of CM production had a significant effect on chemical composition and ruminal protein degradability; (2) determine if CM production plant led to significant differences in chemical composition and ruminal protein degradability; and (3) assess the relative ruminal degradability of protein in CM and SBM.

MATERIALS AND METHODS

Protein Samples

Canola meal samples were collected over 4 yr (2011, 2012, 2013, and 2014), 3 per year, from each of 12 Canadian canola processing plants (total = 144 samples). The 12 plants accounted for the entire CM production in Canada when the studies began; however, a 13th plant has recently come online. About 80% of the CM produced in North America derives from these plants (Carson Callum, Canola Council of Canada, Winnipeg, MB, personal communication). At 11 plants, oil was removed from crushed canola seed by prepress solvent extraction; oil was removed by expeller extraction at

1 plant. The 144 CM samples were identified by plant number (1–12), production year (2011–2014), and replicate within year (1–3). Prior to chemical and in vitro analysis, samples were ground using a laboratory mill fitted with a 1-mm screen (Udy cyclone mill, Udy Corporation, Fort Collins, CO). Three standard proteins were also included in all in vitro incubations: casein (no. C-5890, Sigma Chemicals, St. Louis, MO), solvent-extracted SBM (**SSBM**), and expeller-extracted SBM (**ESBM**). These same standard proteins had been incubated in earlier in vitro studies (Colombini et al., 2011).

Donor Animals and Diets

Ruminal inocula used in the incubations were obtained from 2 lactating Holstein donor cows surgically fitted with ruminal cannulas (Bar Diamond, Parma, ID) and fed a diet composed of 40% alfalfa silage, 20% corn silage, 31.3% ground shelled corn, 8.0% SSBM, 0.4% sodium bicarbonate, and 0.2% salt plus vitamins and trace minerals (on a DM basis) and formulated to 16.5% CP and 1.6 Mcal NE_L/kg DM (at 3× maintenance; NRC, 2001). About 5 min elapsed between collection of inocula and the start of strained ruminal fluid (**SRF**) pre-incubation for incubations conducted in 2011, 2013, and 2014. The University of Wisconsin facility housing donor animals was not available in 2012, necessitating that donor animals be maintained at the US Dairy Forage Research Center farm, which is 40 km from the laboratory. Thus, inocula used in incubations conducted in 2012 were obtained from 2 lactating Holstein donor cows, similarly fitted with ruminal cannulas and fed the same basal diet; however, about 50 min elapsed between collection of inocula and the start of SRF pre-incubations. Surgical care and general maintenance of the animals was as outlined by the guidelines of the University of Wisconsin institutional animal care and use committee.

Chemical Analysis

The CM samples were chemically analyzed in duplicate during the year of collection. Composition data of the 3 standard proteins (casein, SSBM, and ESBM) determined in 2011 were used in computations over all 4 yr. All samples were analyzed for total N (Leco FP-2000 N Analyzer; Leco Instruments, Inc., St. Joseph, MI), DM (method 967.03; AOAC, 1990), ash and OM (method 942.05; AOAC, 1990), sequentially for NDF, ADF, and ADIN using heat stable α -amylase and Na₂SO₃ (Van Soest et al., 1991; Hintz et al., 1996), and for NDIN omitting α -amylase and Na₂SO₃ during

extraction (Licitra et al., 1996). All samples (except casein) were analyzed for buffer insoluble and soluble N using the method of Licitra et al. (1996) with the following modifications: McDougall's buffer (McDougall, 1948) was used for extractions, extracts were filtered through Whatman no. 41 ashless filter paper, insoluble N was determined by combustion (Leco FP-2000) as the difference between the filter plus the retained residue and filter only; the percentage of soluble N was computed as 100 minus insoluble N.

The proportion of total N already degraded at $t = 0$ (FD_0) was originally determined in the year of collection by weighing duplicates of 100 mg of each protein into 50-mL centrifuge tubes and then adding 15 mL of warm (39°C) McDougall's buffer. Samples were swirled and incubated with shaking in the warm room (39°C \pm 0.5) for 2 h. At the end of the buffer-only extraction, 1.25 mL of 65% (wt/vol) trichloroacetic acid (TCA) was added (final concentration 5% wt/vol TCA) and tubes were placed on ice. After 30 min, about 4 mL of sample from each extraction tube was decanted into a 12- by 75-mm plastic tube, and tubes were centrifuged (15 min, 14,000 $\times g$ and 4°C). Supernatants were then transferred to clean tubes and stored at 4°C until analyzed the following day for ammonia, total AA (TAA), and TAA plus oligopeptides as described below. Content of FD_0 in all 144 samples also was reanalyzed in duplicate in 2014 using the procedure just described. The TAA content of proteins ($\mu\text{mol}/\text{mg}$ of N) were determined by hydrolyzing duplicate samples of each protein in 6 N HCl for 24 h at 105°C under a N_2 atmosphere, using a ratio of 1 mg of sample N/5 mL of acid (Block and Weiss, 1956). After hydrolysis, samples were cooled and diluted with distilled water, HCl was removed by vacuum evaporation (Savant SC110 Speedvac Concentrator, Savant Instruments, Inc., Farmingdale, NY), and the residues were redissolved in distilled water and re-evaporated. Residues were then dissolved in 4 mL of 0.10 N HCl and stored (-20°C) for later analysis.

In Vitro Protocol

Incubations were conducted using the Michaelis-Menten inhibitor in vitro method (MMIIV) described by Colombini et al. (2011). All proteins were weighed to the nearest 0.0001 g into 50-mL centrifuge tubes in amounts equivalent to 2.5, 5.0, 10.0, 20.0, and 30.0 mg of N/tube. Whole ruminal contents were collected from beneath the fibrous mat in the rumen and filtered at the barn through 2 layers of cheesecloth into insulated bottles that had flushed previously with CO_2 to remove the air. These filtrates were transported to the laboratory in about 5 min (2011, 2013, 2014) or

about 50 min (2012) and then filtered again through 4 layers of cheesecloth; equal volumes of SRF from each cow were mixed together. To each 1 L of SRF, a solution containing 8 g of maltose (Sigma no. M-2250), 4 g of xylose (Sigma no. X-1500), 4 g of soluble starch (Sigma no. S-2004), and 2.5 g of $NaHCO_3$ dissolved in 150 mL of distilled water, plus a solution containing 4 g of citrus pectin (Sigma no. P-9135) dissolved with heating and slow stirring into 150 mL of McDougall's buffer, were added (total volume 1.30 L). Then, 0.2 mL of the surfactant Antifoam 204 (Sigma no. A-6426) was added, and the inoculum was pre-incubated at 39°C for 3 h under a continuous CO_2 stream. Every hour, pH was measured; if pH was ≤ 6.2 , pH was adjusted to 6.4 by slow addition with stirring of 3 N NaOH. Inhibitor solutions (Broderick, 1987) were prepared by dissolving with stirring 0.267 g of hydrazine sulfate (Sigma no. H-7394) in 35 mL of McDougall's buffer and 0.062 g of chloramphenicol (Sigma no. C-0378) in 35 mL of McDougall's buffer. Addition of these 2 solutions to the 1.30 L of pre-incubated medium yielded 1.37 L of inoculum containing 1.5 mM hydrazine sulfate and 45 mg of chloramphenicol/L. Mercaptoethanol (Sigma no. M-6250) was then added (0.321 mL) as a reducing agent to give 3 mM mercaptoethanol in the inoculum.

While the inoculum was pre-incubating, 5 mL of warm (39°C) McDougall's buffer was added to each incubation tube to suspend all proteins in buffer for 1 h before the start of the incubation. Incubations were begun by dispensing 10 mL of inoculum/tube; immediately after inoculum addition, tube headspace was flushed with CO_2 , and tubes were capped, swirled, and incubated for 2 h at 39°C (± 0.5) in a warm room under continuous mixing with a wrist-arm shaker (set at 100 cycles/min). At the end of the incubation, 1.25 mL of TCA (65% wt/vol) was added to each tube and tubes were placed on ice for 30 min. Next, about 4 mL of sample from each incubation tube was decanted into a 12- by 75-mm plastic tube, and tubes were centrifuged (15 min, 14,000 $\times g$ and 4°C). Supernatants were then transferred to clean tubes, which were stored at 4°C until analyzed the following day as described in the following section. Duplicate blank tubes were included before and after every 40 protein-containing tubes, and incubations typically consisted of 200 to 220 tubes. Incubations of samples from 2011, 2012, and 2013 were replicated 3 times and contained all CM collected in each respective year. Samples collected in 2014, plus those collected in 2013, were run in 6 incubations, with half of the CM from each year included in 3 incubations and the other half included in the other 3 incubations. The 2 replicate observations made for the 2013 CM were averaged to yield 3 rather than 6 incubation observations per CM sample for statistical analysis.

Only the ESBM standard protein was included in the incubations conducted with CM from 2011; all 3 standard proteins were included in incubations conducted with CM collected in 2012, 2013, and 2014.

Determination of Ammonia, TAA, and TAA plus Oligopeptides

Protein hydrolysates and TCA supernatants (from FD₀ extractions and in vitro incubations) were analyzed for ammonia by a phenol-hypochlorite assay adapted to a flow injection analyzer (Lachat Quick-Chem 8000 FIA; Lachat Instruments, Loveland, CO). The *o*-phthalaldehyde (OPA) reaction (Roth, 1971), based on fluorescence from excitation at 340 nm and emission at 450 nm (OPA_F; TAA) or on absorbance at 340 nm (OPA_A; TAA + oligopeptides), was applied to all samples. Those analyses were conducted by flow injection analyzer (Lachat Quick-Chem 8000 FIA) interfaced with a fluorimeter (FS-950 Fluoromat; Kratos Analytical Instruments, Westwood, NJ) for fluorescence (OPA_F assay) and modified with an UV light source (AIS D-1000; Analytical Instrument Systems, Flemington, NJ) for absorbance at 340 nm (OPA_A assay).

The FD₀ values measured by the OPA_A method averaged about 7% of total N, but those determined by the OPA_F method averaged only about 2%; previously, FD₀ values determined for SSBM and ESBM were found to be < 2% by both OPA_A and OPA_F assays (Colombini et al., 2011). This difference was due to a compound (or compounds) in both CM sample extracts and hydrolysates that contributed to A₃₄₀. Initially, A₃₄₀ was measured twice in all buffer extracts, hydrolysates, and in vitro incubations, both with and without OPA reagent in the reaction stream, and the net difference between these values was considered the A₃₄₀ contributed by TAA and oligopeptides. Subsequently, FD₀ values measured using either OPA_A or OPA_F were found to not differ ($P = 0.26$), and data obtained directly by OPA_A were used as obtained (i.e., without correcting for A₃₄₀ detected in the absence of OPA reagent). This approach obviated the need for analyzing samples both with and without OPA reagent.

All computations and nonlinear regression of extents of protein degradation on amount of protein incubated were conducted using the Michaelis-Menten model and methodology described earlier (Broderick and Clayton, 1992; Colombini et al., 2011). Extent of protein degraded (S_D, mg N/mL inoculum) was computed at each level of N addition from net release of N as ammonia plus TAA (and oligopeptides) determined using either the OPA_F or the OPA_A assay with the following equation:

$$S_D = \left\{ \left([\text{NH}_3]_{\text{prot}} - [\text{NH}_3]_{\text{blank}} \right) \times 0.0140067 \right. \\ \left. + \left([\text{TAA}]_{\text{prot}} - [\text{TAA}]_{\text{blank}} \right) / (\text{TAA}/\text{N}) \right\} \times 16.25/10,$$

where [NH₃]_{prot} and [NH₃]_{blank} and [TAA]_{prot} and [TAA]_{blank} are concentrations of ammonia and TAA (μmol/mL) determined by either OPA_A or OPA_F in protein-containing and blank incubation tubes, respectively; 0.0140067 is the mg of N/μmol of ammonia; TAA/N is the ratio of TAA (determined by acid hydrolysis of proteins using either OPA_A or OPA_F) to total N (μmol/mg of N); 16.25 (i.e., 15 + 1.25) is the total tube volume (mL); and 10 is the volume (mL) of added SRF inoculum. Velocity of protein degradation [*v*, mg N/(h · mL inoculum)] was also computed for each protein source at each level of N by dividing the extent of protein degradation (S_D), computed using either TAA assay, by incubation time, *t* (2 h):

$$v = S_D/2.$$

Protein N remaining undegraded at the end of the incubation for each amount of added N (S_{*t*}, mg of N/mL inoculum), was calculated as the difference between the amount of N added at *t* = 0 divided by the volume of inoculum (S₀, mg of N/mL inoculum) and the extent of protein degradation (S_D, defined above), computed using either TAA assay and ammonia analysis:

$$S_t = S_0 - S_D.$$

These data were used to estimate fractional degradation rate (k_d) as the tangent through the origin of the velocity (*v*) versus substrate concentration ([S₀]) curve, and computed from the ratio of maximum velocity (V_{max}) to Michaelis constant (K_m) (i.e., k_d = V_{max}/K_m; Mahler and Cordes, 1966). This was assumed a “blended” or composite ruminal degradation rate contributed by all protein fractions within the sample. This rate was determined using the integrated Michaelis-Menten model (Segal, 1976) as described earlier (Broderick and Clayton, 1992; Colombini et al., 2011):

$$S_t = S_0 - t \times \theta \times K_m + K_m \times \ln(S_0/S_t),$$

where S₀ and S_{*t*} (mg of N/mL inoculum) are as defined above, *t* (h) is the duration of incubation (2 h), and K_m (mg of N/mL of inoculum) is the Michaelis constant defined as that concentration of substrate that gives “half-maximal velocity.” The parameters K_m and θ were estimated by nonlinear regression (SAS, 2013). The θ values were corrected for FD₀ (fraction of total

N already degraded at $t = 0$) to estimate degradation rate, k_d , as follows:

$$k_d = \theta - \text{FD}_0 / t,$$

where $t = 2$ h. The proportion RUP was computed using the model of Waldo et al. (1972) applying the blended rate philosophy and different rates of passage for the insoluble (k_{pi}) and soluble (k_{ps}) N fractions (Columbini et al., 2011):

$$\begin{aligned} \text{RUP} &= \text{Insoluble N} \times k_{pi} / (k_{pi} + k_d) \\ &+ \text{Soluble PrN} \times k_{ps} / (k_{ps} + k_d), \\ \text{RDP} &= 100 - \text{RUP}, \end{aligned}$$

where k_{pi} was set equal to 0.06/h, k_{ps} was set equal to 0.16/h, and soluble PrN = soluble N - FD_0 . As discussed previously, the OPA_A (with correction for A_{340} without OPA reagent) and OPA_F assays yielded FD_0 estimates that were not different. Using either approach to determine FD_0 gave soluble PrN estimates that were not different ($P = 0.94$). Values of soluble PrN determined using the OPA_F assay were applied in all RUP estimates.

Statistical Analysis

Statistical analysis was carried out using the mixed procedures of SAS (2013). The following model was used to assess effects of assaying FD_0 , soluble PrN, k_d , and k_d adjusted to the overall mean k_d , using either the OPA_F or OPA_A assay (corrected for A_{340} determined without OPA reagent):

$$\begin{aligned} Y_{ijkl} &= \mu + Y_i + P_j + M_k + Y \times P_{ij} + Y \times M_{ik} \\ &+ P \times M_{jk} + \text{Rep}_l + E_{ijkl}, \end{aligned}$$

where Y_{ijkl} = dependent variable, μ = overall mean, Y_i = effect of year i ($i = 1$ to 4), P_j = effect of plant j ($j = 1$ to 12), M_k = effect of method k ($k = 1$ to 2, corrected OPA_A or OPA_F), $Y \times P_{ij}$ = interaction of year and plant, $Y \times M_{ik}$ = interaction of year and method, $P \times M_{jk}$ = interaction of plant and method, Rep_l ($l = 1$ to 3 replicates per plant per year), and E_{ijkl} = residual error. All terms were considered fixed, except for Rep_l and E_{ijkl} , which were considered random. The following overall model was used to fit all composition and ruminal degradation data:

$$Y_{ijk} = \mu + Y_i + P_j + Y \times P_{ij} + \text{Rep}_k + \text{Run}_l + E_{ijk},$$

where Y_{ijk} = dependent variable, μ = overall mean, Y_i = effect of year i ($i = 1$ to 4), P_j = effect of plant j ($j = 1$ to 12), $Y \times P_{ij}$ = interaction of year and plant, Rep_k = effect of sample replicate ($k = 1$ to 3 replicates per plant per year), Run_l = effect of in vitro incubation run ($l = 1$ to 3 for 2011, 2012, and 2013, and $l = 6$ for 2014), and E_{ijkl} = residual error. All terms were considered fixed, except for Rep_k , Run_l , and E_{ijkl} , which were considered random. Essentially the same model was used to fit data obtained when CM samples from 2013 and 2014 were incubated in the same in vitro runs except that Y_i = effect of year ($i = 1$ to 2). Sources of variation from sample replicate, in vitro run, year, plant, plus interaction of year and plant were estimated by apportioning total type-3 sums of squares to each source of variation obtained using a SAS GLM version of the overall model. Least squares means estimates are reported; separation of least squares means was conducted at $\alpha = 0.05$ using the PDIF option in the LSMEANS statement. For all statistical analyses, significance was declared at $P \leq 0.05$ and trends at $P \leq 0.10$. Simple linear regressions were used to assess relationships between variables shown in Figures 1 and 2.

RESULTS AND DISCUSSION

Canola Meal Composition

Mean composition data of the CM manufactured over the 4 yr are in Table 1. Significant effects of year of sample collection were detected for most of chemical traits analyzed in this study. Some composition differences, such as DM, organic matter, total N, ADF, insoluble and soluble N, and soluble protein-N, were relatively small but were either significant or highly significant based on year of collection. A surprisingly large difference among years was detected for NDF content, which ranged from about 26% (2011) to almost 31% (2013). However, the overall NDF mean of 28.6% may be compared with the value of 29.8% reported in NRC (2001) tables for "mechanically expelled" CM. Concentrations of most of the macronutrients were also similar to those tabulated by NRC (2001), with CP being the notable exception; the overall mean 6.65% N is equivalent to 41.6% versus 37.8% CP in NRC (2001) tables. Solvent-extracted CM was reported to contain a mean 40.9% CP (Newkirk, 2009). No year effect on NDIN content in CM (expressed as % of total N) was observed and, despite differences in ADIN content, N-fraction B_3 (NDIN - ADIN) did not differ among years.

Concentrations of TAA/N, which were determined by acid hydrolysis, were also different based on the year of sample collection (Table 1). The OPA_F and OPA_A




Figure 1. Regression of ruminal in vitro degradation rate, estimated using *o*-phthaldialdehyde absorbance (OPA_A) methodology, on buffer N-solubility of canola meals collected over 4 yr from 12 production plants.




Figure 2. Regression of ruminal in vitro degradation rate, estimated using *o*-phthaldialdehyde fluorescence (OPA_F) methodology, on buffer N-solubility of canola meals collected over 4 yr from 12 production plants.

values are critical because they are used in the assay to compute the amount of degraded protein from net release of TAA and TAA plus oligopeptides, respectively. Protein AA composition is of course dictated by the cell's DNA template; it seems likely that these apparent differences among years resulted from small "methodological" shifts, such as new standards and standard curves being used over time to compute TAA concentration. However, any effect of changing standard curves would have the same effects on TAA and oligopeptide concentrations determined in the in vitro incubations. Moreover, it was noted that OPA_F and OPA_A values did not differ among processing plant ($P = 0.73$ and 0.74), which allowed mean TAA/N ratios to be applied to all data from all plants within a year. Fraction degraded at $t = 0$ (FD₀) is also important in computing degradation rate. Although an apparent year effect was detected when samples were analyzed by year, when all 144 CM samples were reanalyzed at one time (with the same AA standard curve), no effect of collection year was found ($P = 0.36$).

Mean composition data of the CM manufactured by the 12 crushing plants over the 4 sampling years are in Table 2. Eleven of the 12 crushing plants applied prepress solvent extraction to remove oil; plant 12 was the only facility using the expeller process for oil removal. Expeller processing would be expected to generate more heat and remove less oil, effects that were reflected in meal composition: CM from plant 12 had the highest DM; second highest NDIN, N-fraction

B₃, and insoluble N; numerically highest ADIN; and lowest N concentration. Except for ADIN content and the OPA_F and OPA_A values mentioned earlier, significant or highly significant differences existed among plants for all chemical factors measured in this study. This outcome was probably partly due to the statistical power resulting from the high degree of replication among the processing plants over the 4 sampling years. Of particular interest were the differences in NDIN, N-fraction B₃, and insoluble N, even among CM from plants applying prepress solvent extraction. The relationship of these 3 factors to protein degradation rate and RUP content are discussed in the next section. It was also noted that the 12 crushing plants, although anonymous, were located in 5 Canadian provinces (Quebec, Ontario, Manitoba, Saskatchewan, and Alberta). Thus, the canola seed processed by each plant originated from various canola cultivars grown over widely varying geographical areas with a range of soil types and environmental conditions. Mean temperature and moisture had greater effects than cultivar on oil and protein content of canola seed grown in Australia from 1985 through 1994 (Pritchard et al., 2000). Canola cultivar had greater effects on fatty acid composition and spring temperatures and moisture had greater effects on oil and protein content of canola seed grown in Canadian trials (McCartney et al., 2004). Timing and amount of N fertilization interacted with soil type to influence composition and yield of canola seed (Brennan, 2016).

Table 1. Mean chemical composition of canola meals from all 4 yr

| Item | Production year | | | | SE | Probability | | |
|---|--------------------|-------------------|-------------------|--------------------|-------|-------------|--------|--------------|
| | 2011 | 2012 | 2013 | 2014 | | Year | Plant | Year × plant |
| DM, % | 90.6 ^{ab} | 90.8 ^a | 90.9 ^a | 90.4 ^b | 0.14 | 0.039 | <0.001 | <0.001 |
| OM, % of DM | 93.7 ^a | 91.8 ^c | 91.9 ^c | 92.4 ^b | 0.11 | <0.001 | 0.051 | 0.719 |
| Total N, % of DM | 6.67 ^b | 6.67 ^b | 6.79 ^a | 6.50 ^c | 0.019 | <0.001 | <0.001 | <0.001 |
| NDF, % of DM | 25.9 ^c | 27.8 ^b | 30.6 ^a | 30.0 ^a | 0.32 | <0.001 | <0.001 | <0.001 |
| ADF, % of DM | 18.1 ^b | 19.2 ^a | 18.0 ^b | 18.9 ^a | 0.18 | <0.001 | 0.012 | 0.002 |
| NDIN, % of total N | 21.0 | 22.5 | 20.9 | 20.5 | 0.98 | 0.443 | 0.013 | 0.071 |
| ADIN, % of total N | 7.1 ^a | 7.6 ^a | 4.1 ^c | 6.2 ^b | 0.21 | <0.001 | 0.539 | 0.355 |
| Fraction B ₃ , ¹ % of total N | 13.9 | 14.8 | 16.8 | 14.3 | 0.97 | 0.123 | 0.045 | 0.215 |
| Insoluble N, % of total N | 74.5 ^a | 71.2 ^b | 71.5 ^b | 72.7 ^{ab} | 0.65 | 0.002 | <0.001 | 0.028 |
| Soluble N, % of total N | 25.5 ^b | 28.8 ^a | 28.6 ^a | 27.3 ^{ab} | 0.65 | 0.002 | <0.001 | 0.028 |
| Soluble protein N, ² % of total N | 23.4 ^b | 26.7 ^a | 26.4 ^a | 25.2 ^{ab} | 0.65 | 0.002 | <0.001 | 0.029 |
| FD ₀ , ³ % of total N | 2.13 | 2.15 | 2.15 | 2.14 | 0.041 | 0.356 | <0.001 | 0.607 |
| OPA _A , ⁴ μmol/mg of N | 43.3 ^b | 44.1 ^a | 42.9 ^b | 43.2 ^b | 0.28 | 0.015 | 0.734 | 0.994 |
| OPA _F , ⁵ μmol/mg of N | 30.2 ^c | 31.3 ^b | 35.3 ^a | 35.8 ^a | 0.24 | <0.001 | 0.743 | 0.984 |

^{a-c}LSM with different superscript letters among years are different ($P < 0.05$).

¹Fraction B₃ = NDIN - ADIN.

²Soluble protein N = Soluble N - FD₀.

³FD₀ = fraction total N present as degraded N (ammonia + free AA) at $t = 0$.

⁴OPA (*o*-phthalaldehyde) color factor (A₃₄₀) in μmol of Leu equivalents/mg of N.

⁵OPA fluorescence factor in μmol of Leu equivalents/mg of N.

Table 2. Mean chemical compositions of canola meals from all 12 processing plants

| Item | Processing plant | | | | | | | | | | | | Probability | | |
|---|---------------------|----------------------|--------------------|---------------------|---------------------|----------------------|--------------------|---------------------|---------------------|--------------------|---------------------|----------------------|-------------|--------|--------------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | SE | Plant | Year × plant |
| Processing method ¹ | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Solv. | Exp. | | | |
| DM, % | 89.9 ^{de} | 89.5 ^e | 89.5 ^e | 89.6 ^e | 89.8 ^e | 90.5 ^{cd} | 90.8 ^{bc} | 90.9 ^{bc} | 91.3 ^b | 90.4 ^{cd} | 91.0 ^{bc} | 94.9 ^a | 0.22 | <0.001 | <0.001 |
| OM, % of DM | 92.3 ^{abc} | 92.3 ^{abc} | 92.2 ^{bc} | 92.7 ^a | 92.4 ^{abc} | 92.0 ^c | 92.7 ^a | 92.7 ^a | 92.4 ^{abc} | 92.5 ^{ab} | 92.6 ^{ab} | 92.6 ^{ab} | 0.17 | 0.051 | 0.719 |
| Total N, % of DM | 6.85 ^a | 6.59 ^d | 6.64 ^{cd} | 6.85 ^a | 6.72 ^{bc} | 6.78 ^{ab} | 6.50 ^e | 6.69 ^c | 6.65 ^c | 6.86 ^a | 6.58 ^{de} | 6.15 ^f | 0.030 | <0.001 | <0.001 |
| NDF, % of DM | 26.3 ^e | 29.3 ^{bc} | 27.8 ^{cd} | 27.4 ^{de} | 28.2 ^{cd} | 28.1 ^{cd} | 28.4 ^{cd} | 30.3 ^{ab} | 31.4 ^a | 27.7 ^{de} | 28.0 ^{cd} | 30.6 ^{ab} | 0.55 | <0.001 | <0.001 |
| ADF, % of DM | 18.1 ^{cd} | 18.7 ^{abcd} | 19.0 ^{ab} | 17.9 ^d | 18.2 ^{bcd} | 18.1 ^{cd} | 19.1 ^a | 19.2 ^a | 18.8 ^{abc} | 18.0 ^{cd} | 19.0 ^{ab} | 18.4 ^{abcd} | 0.31 | 0.012 | 0.002 |
| NDIN, % of total N | 18.3 ^d | 22.7 ^{abc} | 21.0 ^{cd} | 20.4 ^{bcd} | 19.7 ^{cd} | 22.3 ^{abcd} | 19.4 ^{cd} | 21.3 ^{bcd} | 26.3 ^a | 19.6 ^{cd} | 19.1 ^{cd} | 24.6 ^{ab} | 1.61 | 0.013 | 0.071 |
| ADIN, % of total N | 5.9 | 6.7 | 6.4 | 5.9 | 6.0 | 6.1 | 6.6 | 6.5 | 6.4 | 5.9 | 6.2 | 6.9 | 0.37 | 0.539 | 0.355 |
| Fraction B ₃ , ² % of total N | 12.4 ^c | 16.0 ^{abc} | 14.7 ^{bc} | 14.5 ^{bc} | 13.7 ^{bc} | 16.2 ^{abc} | 12.8 ^c | 14.8 ^{bc} | 19.9 ^a | 13.7 ^{bc} | 12.9 ^c | 17.7 ^{ab} | 1.62 | 0.045 | 0.215 |
| Insoluble N, % of total N | 62.5 ^f | 77.8 ^{ab} | 74.7 ^{bc} | 75.6 ^b | 79.7 ^a | 71.3 ^{de} | 71.8 ^{cd} | 69.1 ^{de} | 68.4 ^e | 68.2 ^e | 71.1 ^{de} | 79.3 ^a | 1.12 | <0.001 | 0.028 |
| Soluble N, % of total N | 37.5 ^a | 22.2 ^{df} | 25.3 ^{de} | 24.4 ^e | 20.3 ^f | 28.7 ^{bc} | 28.2 ^{cd} | 30.9 ^{bc} | 31.6 ^b | 31.8 ^b | 28.9 ^{bc} | 20.7 ^f | 1.12 | <0.001 | 0.028 |
| FD ₀ , ³ total N | 2.22 ^a | 2.12 ^{cd} | 2.16 ^{bc} | 2.11 ^d | 2.10 ^d | 2.17 ^b | 2.19 ^{ab} | 2.18 ^b | 2.17 ^b | 2.13 ^{cd} | 2.10 ^d | 2.06 ^e | 0.043 | <0.001 | 0.607 |
| Soluble protein N, ⁴ % of total N | 35.3 ^a | 20.1 ^{fg} | 23.2 ^{df} | 22.3 ^f | 18.2 ^g | 26.5 ^{cd} | 26.0 ^{de} | 28.7 ^{bcd} | 29.5 ^{bc} | 29.7 ^b | 26.8 ^{bcd} | 18.7 ^g | 1.12 | <0.001 | 0.029 |
| FD ₀ , ⁴ % of total N | 2.43 ^a | 2.03 ^{de} | 2.15 ^c | 2.00 ^e | 1.93 ^{ef} | 2.18 ^{bc} | 2.27 ^b | 2.23 ^{bc} | 2.13 ^{cd} | 2.30 ^b | 2.19 ^{bc} | 1.88 ^f | 0.040 | <0.001 | <0.001 |
| OPA ₅ , ⁵ μmol/mg of N | 43.4 | 43.9 | 43.2 | 43.3 | 43.1 | 44.1 | 43.1 | 43.4 | 43.7 | 43.8 | 43.1 | 42.7 | 0.48 | 0.734 | 0.994 |
| OPA _F , ⁶ μmol/mg of N | 33.3 | 33.5 | 32.9 | 33.2 | 32.8 | 33.8 | 32.8 | 33.1 | 33.6 | 33.3 | 32.9 | 32.7 | 0.42 | 0.743 | 0.984 |

^{a-f}LSM with different superscript letters among plants are different ($P < 0.05$).

¹Canola meal was processed to remove oil by either solvent extraction (Solv.) or expeller extraction (Exp.).

²Fraction B₃ = NDIN - ADIN.

³FD₀ = fraction total degraded N present at $t = 0$, in the form of ammonia plus free AA [determined using o-phthalaldehyde (OPA) fluorescence method].

⁴Soluble protein N = Soluble N - FD₀.

⁵OPA absorbance factor (A₃₄₀) in μmol of Leu equivalents/mg of N.

⁶OPA fluorescence factor in μmol of Leu equivalents/mg of N.

Determination of Ruminant Protein Degradability

Degradation data obtained for the standard proteins casein, SSBM, and ESBM in the MMIIV system are in Table 3. Michaelis constant (K_m) and maximum velocity (V_{max}) are reported in addition to degradation rate and proportions of RUP and RDP. Values of K_m represent affinity of the substrate for the enzyme system (Mahler and Cordes, 1966) and are inversely related to reaction rate. The maximal rate of an enzymatic process is represented by V_{max} (Segal, 1976), which should be similar for all substrates tested; V_{max} was not different among the 3 standard proteins, indicating that the MMIIV system met this general requirement. As expected, detection of oligopeptides in addition to TAA yielded higher k_d and lower RUP estimates for the OPA_A method versus the OPA_F method. This outcome was observed previously for these same proteins (Colombini et al., 2011). Ruminant metabolism of small peptides is very dynamic (Chen et al., 1987; Broderick et al., 1988), and peptides contribute small but significant amounts to the total AA flow from the rumen (Choi et al., 2003). When tested earlier, the standard proteins yielded somewhat different degradation rates and RUP values (Colombini et al., 2011): RUP estimates made using OPA_F and OPA_A methodology were, respectively, 21 and 27% (casein), 35 and 38% (SSBM), and 42 and 48% (ESBM). Values of 31 and 35% RUP were observed for SSBM in the present study; however, 54 and 59% RUP were obtained for ESBM. The difference between trials for ESBM was partly due to greater N-solubility in the current study (8.5%) versus the lower earlier measurement on the same protein (4.5%). Applying the same blended rate to proteins of greater solubility results in a greater RUP estimate because of the higher passage rate from the rumen. One may speculate that the ideal high RUP protein would be both resistant to ruminal degradation but soluble and thus passing rapidly with the liquid phase. Degradation rates and RUP values (at DMI = 4% of BW) for SSBM and ESBM assigned in NRC (2001) tables are, respectively, 0.075 and 0.024/h and 43 and 69%. Although based on very different *in situ* methods, the SSBM/ESBM ratios for the NRC (2001) are 3.1 (degradation rate) and 0.6 (RUP), similar to the values observed for these ratios in the present study. These results indicated that the MMIIV system yielded reliable relative estimates of protein degradation rate and RUP values for the standard proteins and would thus be expected to yield reliable protein degradability data for the CM sample set. Although θ and thus k_d were similar among years, RUP values estimated by both the OPA_A and OPA_F assays differed by incubation year (Table 3). This outcome likely results from applying

constant proportions of soluble and insoluble protein for each standard protein plus constant passage rates; this approach yields RUP values with lower variation than is obtained for k_d .

In addition to N composition data, mean ruminal protein degradation results for the CM manufactured over the 4 sampling years are in Table 4. Large differences existed among years in θ ; this result indicated that large differences were present in degradation rate k_d , which is computed by discounting θ for FD_0 (fraction of total N already degraded at $t = 0$). Lowest mean θ was observed in 2012, when the SRF had to be transported about 40 km for 50 min before pre-incubation. However, a year effect was not observed for θ and k_d determined for the 3 standard proteins (Table 3). Earlier, substantial day-to-day differences in degradation rate also were observed for the same proteins incubated with ruminal inocula prepared from SRF within 5 min of collection from the same donor cows and fed the same diets (Broderick et al., 2004). Therefore, we decided to directly test the effect of year by rerunning CM collected in 2013 in the same incubations as CM collected in 2014 to assess whether CM protein degradability varied because of production year; these data are in Table 5. As shown in Table 1, N composition also differed for CM from 2013 and 2014. However, no differences existed between years ($P \geq 0.36$) for θ , k_d , RUP, and RDP. Thus, apparent differences among years for the degradation parameters shown in Table 4 reflected the high variability in degradative activity among ruminal inocula. We decided to adjust θ observed for each protein within each incubation, by dividing the observed θ by the ratio: mean incubation θ /overall 4-year mean θ . The adjusted θ values, plus the k_d , RUP, and RDP computed from these values, are reported in Table 4 and again show no differences in CM degradability among years of sample collection. Mean k_d and RUP values for CM for both assays over all 4 yr were 0.09/h and 45%, compared with k_d and RUP of 0.16/h and 33% for SSBM and 0.05/h and 56% for ESBM. Maxin et al. (2013a) reported *in situ* k_d and RUP for CM of 0.06/h and 53% versus k_d and RUP of 0.09/h and 42% for SSBM. Note that, unlike in the *in situ* method, soluble protein contributes to the measured degradation rate and RUP in the MMIIV system.

Mean ruminal protein degradation results using adjusted θ values for CM produced by the 12 canola crushing plants over all 4 sampling years are in Table 6. These results were highly significantly different among plants. Note that for both OPA_A and OPA_F data the magnitude of θ and adjusted θ values are almost identical. However, the spread of mean separations was somewhat narrower after adjustment: θ mean separations were a to g, but adjusted θ mean separations were a to f

(OPA_A); θ mean separations were a to f, but adjusted θ mean separations were a to d (OPA_F). What was striking was the wide differences in degradability among CM produced by the 12 plants that were consistent between the 2 methods: k_d ranged from 0.069 to 0.113/h and RUP from 51 to 43% (OPA_A), and k_d ranged from 0.075 to 0.120/h and RUP from 49 to 41% (OPA_F). The CM produced by plants 5 (a prepress solvent extraction plant) and 12 (the sole expeller plant) averaged 7.7 and 7.2 percentage units greater RUP than CM produced by prepress solvent extraction plants 1 and 10, a mean 18% greater RUP by both methods.

The similarity of CM k_d and RUP values determined by the OPA_A and OPA_F methods was surprising and unexpected. Research with casein, ESBM, and SSBM conducted previously (Colombini et al., 2011) indicated more rapid k_d and lower RUP estimates using the OPA_A assay, reflecting the contribution of oligopeptide release with OPA_A assay. In the present study, use of the OPA_A assay yielded k_d that were 15% greater for casein and 25% greater for the 2 SBM, while RUP estimates were reduced an average 11% across the 3 standard proteins (Table 3). Analysis of CM data before

adjustment for run variation yielded mean k_d of 0.095 and 0.092/h for, respectively, the OPA_A and OPA_F assays ($P = 0.039$); however, this difference disappeared after the adjustment. Computation of k_d derives in part from dividing net TAA (OPA_F) or TAA plus peptide (OPA_A) release by the TAA/N constants obtained from HCl hydrolysis; overall mean TAA/N constants were 43.4 and 33.2 $\mu\text{mol}/\text{mg N}$ for, respectively, the OPA_A and OPA_F assays (Table 1). These results suggest that peptide accumulation makes a smaller contribution to the degradation rates estimated for CM protein versus the proteins in casein and SBM. Substantial differences in rates of disappearance of various di- and tripeptides have been observed (Broderick et al., 1988); it may be that AA composition of the peptides released during ruminal CM degradation are such that they are rapidly catabolized by ruminal microorganisms.

Clearly, consistent differences existed among processing plants in the CM that they produced. Assuming intestinal digestibility was not impaired, RUP differences of the magnitudes estimated by the MMIV procedure imply that MP and AA supply could differ substantially among CM produced at different plants.

Table 3. Composition and ruminal in vitro degradation results for casein, solvent soybean meal (SSBM), and expeller soybean meal (ESBM) included in the incubations

| Item | Standard protein | | | SE | Probability | | | SSBM: ESBM |
|--|------------------|-------|-------|--------|-------------|-------|-----------------------|---------------|
| | Casein | SSBM | ESBM | | Protein | Year | Year \times protein | |
| Composition | | | | | | | | |
| Total N, % of DM | 16.11 | 8.44 | 8.14 | — | — | — | — | — |
| Insoluble N, % of total N | 0 | 80.1 | 91.5 | — | — | — | — | — |
| Soluble N, % of total N | 100.0 | 19.9 | 8.5 | — | — | — | — | — |
| FD ₀ , ¹ % of total N | 0.25 | 1.09 | 1.27 | — | — | — | — | — |
| Soluble protein N, ² % of total N | 99.7 | 18.9 | 7.2 | — | — | — | — | — |
| OPA _A results ³ | | | | | | | | |
| θ , per h | 0.818 | 0.183 | 0.062 | 0.1555 | 0.002 | 0.709 | 0.691 | — |
| K_m , ⁴ mg of N/mL of SRF | 0.197 | 0.857 | 3.283 | 0.2143 | <0.001 | 0.388 | 0.006 | — |
| V_{max} , ⁵ mg of N/h per mL of SRF | 0.161 | 0.145 | 0.198 | 0.0222 | 0.712 | 0.646 | 0.202 | — |
| k_d , ⁶ per h | 0.816 | 0.177 | 0.055 | 0.1554 | 0.002 | 0.708 | 0.692 | 3.23 |
| RUP, ⁷ % of total N | 16.5 | 30.5 | 53.6 | 2.99 | <0.001 | 0.024 | 0.032 | 0.57 |
| RDP, ⁷ % of total N | 83.5 | 69.5 | 46.4 | 2.99 | <0.001 | 0.024 | 0.032 | 1.50 |
| OPA _F results ⁸ | | | | | | | | |
| θ , per h | 0.713 | 0.149 | 0.049 | 0.0481 | <0.001 | 0.160 | 0.228 | — |
| K_m , ⁴ mg of N/mL of SRF | 0.135 | 0.736 | 2.701 | 0.1197 | <0.001 | 0.054 | 0.001 | — |
| V_{max} , ⁵ mg of N/h per mL of SRF | 0.096 | 0.101 | 0.132 | 0.0088 | 0.233 | 0.002 | 0.267 | — |
| k_d , ⁶ per h | 0.712 | 0.145 | 0.044 | 0.0480 | <0.001 | 0.159 | 0.227 | 3.31 |
| RUP, ⁷ % of total N | 18.5 | 34.9 | 59.0 | 3.84 | <0.001 | 0.007 | 0.093 | 0.59 |
| RDP, ⁷ % of total N | 81.5 | 65.1 | 41.0 | 3.84 | <0.001 | 0.007 | 0.093 | 1.59 |

¹FD₀ = fraction total N present as degraded N (ammonia and free AA) at $t = 0$. Values in this table are FD₀ means determined using OPA_F and OPA_A methodology.

²Soluble protein N = Soluble N – FD₀.

³Degradation determined from net release of ammonia plus α -phthalaldehyde (OPA) absorbance (A_{340}), which includes free AA plus oligopeptides.

⁴Michaelis constant per mL of strained ruminal fluid (SRF) in inoculum.

⁵Maximum velocity per mL of strained ruminal fluid (SRF) in inoculum.

⁶Degradation rate (k_d) = $\theta - (\text{FD}_0/100)/\text{incubation time (2 h)}$.

⁷RUP = insoluble N \times 0.06/(0.06 + k_d) + soluble protein N \times 0.16/(0.16 + k_d); RDP = 100 – RUP.

⁸Degradation determined from net release of ammonia plus OPA fluorescence, which includes only free AA.

Newkirk et al. (2003) found significant differences in digestibility in broiler chickens of CP and 16 AA due to plant of origin for 31 CM samples obtained from 7 crushing plants, all of which applied prepress solvent extraction; effects were greatest for Lys, the digestibility of which ranged from 70.4 to 82.1% among CM from the 7 plants. The authors attributed this effect to the Maillard reaction during toasting, which is used to remove hexane (Classen et al., 2004); these findings were later confirmed in swine (Thacker and Newkirk, 2005). The differences we observed in ruminal protein degradation in CM produced by the various crushing plants could be explained by differences in degree of heating when meal is desolventized (Moshtaghi Nia and Ingalls, 1995). Heat-treating protein concentrates to improve RUP requires finding the optimum balance between ruminal protein escape and intestinal protein digestion (e.g., Faldet et al., 1992). Huhtanen et al. (2011) observed numerically greater milk yield response to heat-treated CM versus conventional CM, although this difference was not significant ($P = 0.30$). We suggest that application of consistent temperature treatments during prepress solvent processing among canola crushing plants would result in production of

CM with more uniform RUP content and nutritional value.

Assessing plant \times year interactions was not an original objective and was considered beyond the scope of this research; however, this interaction proved to be highly significant for several composition and ruminal degradation traits in this data set (Table 6). This finding may have partly resulted from the strong statistical power of this study referred to earlier. We estimated the relative contributions to total variance by apportioning GLM type-3 sums of squares for the different model components (Table 7). The statistical models explained an estimated 41 to 83% of observed variation. Except for the fraction degraded at $t = 0$ (for both FD_{0A} and FD_{0F}), sample replicate contributed little to overall variance. Incubation run and year were also unimportant except for unadjusted θ . Averaging OPA_A and OPA_F data, plant accounted for 35% (k_d) and 23% (RUP) of total variance, and plant \times year accounted for 15% (k_d) and 18% (RUP) of total variance, suggesting that plant was substantially more important as a factor determining rate of CM protein degradation.

We used simple regressions of degradation rate k_d , determined by both OPA_A and OPA_F methodology, on

Table 4. Mean yearly ruminal in vitro degradation results for canola meals collected over 4 production years

| Item | Production year | | | | SE | Probability | |
|--|--------------------|--------------------|--------------------|--------------------|--------|-------------|---------------------|
| | 2011 | 2012 | 2013 | 2014 | | Year | Year \times plant |
| Composition, % of total N | | | | | | | |
| Insoluble N | 74.5 ^a | 71.2 ^b | 71.5 ^b | 72.7 ^{ab} | 0.65 | 0.002 | 0.028 |
| Soluble N | 25.5 ^b | 28.8 ^a | 28.6 ^a | 27.3 ^{ab} | 0.65 | 0.002 | 0.028 |
| FD_0 ¹ | 2.13 | 2.15 | 2.15 | 2.14 | 0.041 | 0.356 | 0.607 |
| Soluble protein N ² | 23.1 ^c | 27.1 ^a | 26.3 ^{ab} | 25.2 ^b | 0.65 | <0.001 | 0.034 |
| OPA _A results ³ | | | | | | | |
| FD_{0A} , ⁴ % of total N | 7.07 ^b | 7.20 ^a | 7.02 ^b | 7.18 ^a | 0.105 | <0.001 | 0.003 |
| θ , per h | 0.113 ^c | 0.094 ^d | 0.142 ^b | 0.149 ^a | 0.003 | <0.001 | <0.001 |
| Adjusted θ , ⁵ per h | 0.125 | 0.125 | 0.125 | 0.125 | 0.0018 | 0.997 | <0.001 |
| k_d , ⁶ per h | 0.090 | 0.089 | 0.090 | 0.090 | 0.0018 | 0.954 | <0.001 |
| RUP, ⁷ % of total N | 46.0 | 47.1 | 45.8 | 46.1 | 0.49 | 0.199 | <0.001 |
| RDP, ⁷ % of total N | 54.0 | 52.9 | 54.2 | 53.9 | 0.49 | 0.199 | <0.001 |
| OPA _F results ⁸ | | | | | | | |
| FD_{0F} , ⁹ % of total N | 2.13 | 2.15 | 2.15 | 2.14 | 0.041 | 0.356 | 0.607 |
| θ , per h | 0.100 ^c | 0.078 ^d | 0.122 ^b | 0.127 ^a | 0.0029 | <0.001 | <0.001 |
| Adjusted θ , ⁵ per h | 0.107 | 0.107 | 0.107 | 0.107 | 0.0017 | 1.000 | <0.001 |
| k_d , ⁶ per h | 0.094 | 0.098 | 0.095 | 0.096 | 0.0017 | 0.382 | <0.001 |
| RUP, ⁷ % of total N | 44.4 | 44.7 | 44.3 | 44.1 | 0.41 | 0.738 | <0.001 |
| RDP, ⁷ % of total N | 55.6 | 55.3 | 55.7 | 55.9 | 0.41 | 0.738 | <0.001 |

^{a-d}LSM with different superscript letters among years are different ($P < 0.05$).

¹ FD_0 = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$, where OPA = *o*-phthalaldehyde.

²Soluble protein N = Soluble N - FD_0 .

³Degradation determined from net release of ammonia plus OPA absorbance (A_{340}), which includes free AA plus oligopeptides.

⁴ FD_{0A} = fraction total N present as degraded N [ammonia and free AA plus oligopeptides (A_{340})] at $t = 0$.

⁵Adjusted θ computed by dividing θ observed for individual proteins in each incubation by the ratio: mean incubation θ /overall mean θ .

⁶Degradation rate (k_d) = adjusted θ - ($FD_0/100$)/incubation time (2 h).

⁷RUP = insoluble N \times 0.06/(0.06 + k_d) + soluble protein N \times 0.16/(0.16 + k_d); RDP = 100 - RUP.

⁸Degradation determined from net release of ammonia plus OPA fluorescence, which includes only free AA.

⁹ FD_{0F} = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$.

Table 5. Ruminal in vitro degradation results for canola meals collected in 2013 and 2014 and incubated simultaneously in the same incubations

| Item | Year | | | Probability | | |
|--|-------|-------|--------|-------------|--------|--------------|
| | 2013 | 2014 | SE | Year | Plant | Year × plant |
| Composition, % of total N | | | | | | |
| Insoluble N | 71.5 | 72.7 | 0.39 | 0.032 | <0.001 | 0.057 |
| Soluble N | 28.6 | 27.3 | 0.39 | 0.032 | <0.001 | 0.057 |
| FD ₀ ¹ | 2.15 | 2.14 | 0.047 | 0.275 | <0.001 | 0.987 |
| Soluble protein N ² | 26.4 | 25.2 | 0.39 | 0.033 | <0.001 | 0.058 |
| OPA _A results ³ | | | | | | |
| FD _{0A} , ⁴ % of total N | 7.02 | 7.18 | 0.11 | 0.001 | <0.001 | 0.467 |
| θ, per h | 0.148 | 0.150 | 0.0076 | 0.547 | <0.001 | <0.001 |
| k _d , ⁵ per h | 0.113 | 0.114 | 0.0076 | 0.853 | <0.001 | <0.001 |
| RUP, ⁶ % of total N | 41.0 | 40.8 | 1.57 | 0.613 | <0.001 | <0.001 |
| RDP, ⁶ % of total N | 59.0 | 59.2 | 1.57 | 0.613 | <0.001 | <0.001 |
| OPA _F results ⁷ | | | | | | |
| FD _{0F} , ⁸ % of total N | 2.15 | 2.14 | 0.047 | 0.275 | <0.001 | 0.987 |
| θ, per h | 0.126 | 0.128 | 0.0062 | 0.362 | <0.001 | <0.001 |
| k _d , ⁵ per h | 0.115 | 0.117 | 0.0063 | 0.145 | <0.001 | <0.001 |
| RUP, ⁶ % of total N | 40.5 | 39.8 | 1.29 | 0.065 | <0.001 | <0.001 |
| RDP, ⁶ % of total N | 59.5 | 60.2 | 1.29 | 0.065 | <0.001 | <0.001 |

¹FD₀ = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$, where OPA = *o*-phthalaldehyde.

²Soluble protein N = Soluble N – FD₀.

³Degradation determined from net release of ammonia plus OPA absorbance (A₃₄₀), which includes free AA plus oligopeptides.

⁴FD_{0A} = fraction total N present as degraded N [ammonia and free AA plus oligopeptides (A₃₄₀)] at $t = 0$.

⁵Degradation rate (k_d) = $\theta - (\text{FD}_0/100)/\text{incubation time (2 h)}$.

⁶RUP = insoluble N × 0.06/(0.06 + k_d) + soluble protein N × 0.16/(0.16 + k_d); RDP = 100 – RUP.

⁷Degradation determined from net release of ammonia plus OPA fluorescence, which includes only free AA.

⁸FD_{0F} = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$.

soluble N, NDIN, and N-fraction B₃ (as proportions of total N) to assess their relationships to CM protein degradability. Regressions on NDIN and fraction B₃ yielded correlation coefficients (R²) of only 0.021 and 0.015, respectively. However, regressions on soluble N yielded R² of 0.402 for k_d determined by OPA_A (Figure 1) and 0.422 for k_d determined by OPA_F (Figure 2). Of course, regressing k_d on insoluble N yielded identical R² but negative slopes. Although about 60% of the variation in k_d was still unexplained, this finding was of interest. Heat treatment reduces protein solubility in cottonseed meal (Craig and Broderick, 1981), soybeans (Faldet and Satter, 1991), SBM (Borucki Castro et al., 2007), CM (Moshtaghi Nia and Ingalls, 1995), and other proteins. Heat treatment of protein also promotes the Maillard reaction between lysine residues and reducing sugars present in feeds that alter susceptibility of proteins to digestion in the rumen (Faldet et al., 1992) and small intestine (Friedman, 1996). Extent of the Maillard reaction is directly related to extent of ruminal protein escape (Boucher et al., 2009), and protein solubility/insolubility could thus be correlated to several factors influencing ruminal protein degradation and escape.

Substantial in vivo evidence exists for differences in lactation performance between dairy cows fed SSBM and ESBM (e.g., Broderick et al., 1990) and between dairy

cows fed SSBM and CM (Huhtanen et al., 2011; Martineau et al., 2013). Greater milk and protein yield were observed when equal CP diets containing CM replaced those containing SSBM; lower ruminal concentrations of ammonia and branched-chain VFA suggested that at least part of the positive effect of replacing SSBM was due to greater RUP on the CM diets (Broderick et al., 2015). However, we know of no published in vivo data showing differences in milk or protein yield among CM coming from different processing plants. Using omasal sampling, Brito et al. (2007) found a numeric difference in RUP flow of 90 g/d when CM substituted for equal CP from SSBM in cows consuming about 1.6 kg CP from either supplement. Mean RUP estimated by both OPA_A and OPA_F methods was 33% for SSBM and 45% over all CM plants from all 4 sampling years; if these RUP values were applied in vivo, then RUP flow should have been about 180 g/d greater on CM than SSBM, a result about twice that observed in the trial of Brito et al. (2007). Maxin et al. (2013b) determined differences in whole-body irreversible loss of Lys, Met, and His (estimates of net AA absorption) in dairy cows fed SBM and CM; Lys irreversible loss was higher and irreversible loss of Met and His was numerically higher on CM versus SBM. Greater contribution of MP from CM than SSBM is consistent with our present in vitro findings and emphasizes the importance of improving

Table 6. Mean ruminal in vitro degradation data for canola meals produced at all 12 processing plants over all 4 yr

| Item | Processing plant | | | | | | | | | | | | Probability | | |
|--|--------------------|---------------------|----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|----------------------|---------------------|---------------------|--------------------|-------------|--------|-----------------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | SE | Plant | Year × plant |
| Composition, % of total N ¹ | | | | | | | | | | | | | | | |
| Insoluble N | 62.5 ^f | 77.8 ^{ab} | 74.7 ^{bc} | 75.6 ^b | 79.7 ^a | 71.3 ^{de} | 71.8 ^{cd} | 69.1 ^{de} | 68.4 ^c | 68.2 ^c | 71.1 ^{de} | 79.3 ^a | 1.12 | <0.001 | 0.028 |
| Soluble N | 37.5 ^a | 22.2 ^{ef} | 25.3 ^{de} | 24.4 ^c | 20.3 ^f | 28.7 ^{bc} | 28.2 ^{cd} | 30.9 ^{bc} | 31.6 ^b | 31.8 ^b | 28.9 ^{bc} | 20.7 ^f | 1.12 | <0.001 | 0.028 |
| FD ₀ ² | 2.22 ^a | 2.12 ^{cd} | 2.16 ^{bc} | 2.11 ^d | 2.10 ^d | 2.17 ^b | 2.19 ^{ab} | 2.18 ^b | 2.17 ^b | 2.13 ^{cd} | 2.10 ^d | 2.06 ^c | 0.043 | <0.001 | 0.607 |
| Soluble protein N ³ | 35.3 ^a | 20.1 ^{fg} | 23.2 ^{ef} | 22.3 ^f | 18.2 ^g | 26.5 ^{cd} | 26.0 ^{de} | 28.7 ^{bcd} | 29.5 ^{bc} | 29.7 ^b | 26.8 ^{bcd} | 18.7 ^g | 1.12 | <0.001 | 0.029 |
| OPA _A results ⁴ | | | | | | | | | | | | | | | |
| FD _{0A} , % of total N | 7.50 ^a | 6.87 ^{fg} | 7.24 ^{cd} | 6.86 ^{gh} | 6.73 ^h | 7.09 ^e | 7.43 ^{ab} | 7.27 ^c | 7.11 ^{de} | 7.30 ^{bc} | 7.01 ^{ef} | 7.01 ^{ef} | 0.112 | <0.001 | 0.003 |
| θ, per h | 0.149 ^a | 0.111 ^{fg} | 0.127 ^{bcd} | 0.114 ^{ef} | 0.104 ^g | 0.124 ^{cd} | 0.133 ^b | 0.132 ^{bc} | 0.122 ^{de} | 0.141 ^a | 0.132 ^{bc} | 0.104 ^g | 0.0041 | <0.001 | <0.001 |
| Adjusted θ, per h | 0.151 ^a | 0.112 ^{ef} | 0.127 ^{bc} | 0.116 ^{de} | 0.104 ^f | 0.123 ^{cd} | 0.134 ^b | 0.132 ^b | 0.121 ^{cd} | 0.145 ^a | 0.134 ^b | 0.104 ^f | 0.0031 | <0.001 | <0.001 |
| k _d , per h | 0.113 ^a | 0.078 ^{ef} | 0.091 ^{bc} | 0.081 ^{de} | 0.071 ^{fg} | 0.088 ^{cd} | 0.097 ^b | 0.096 ^b | 0.086 ^{cde} | 0.108 ^a | 0.099 ^b | 0.069 ^g | 0.0031 | <0.001 | <0.001 |
| RUP, % of total N | 43.2 ^{gh} | 47.6 ^c | 45.4 ^{def} | 47.3 ^{cd} | 50.1 ^{ab} | 46.6 ^{cde} | 44.2 ^{fgh} | 45.1 ^{de} | 48.1 ^{bc} | 42.5 ^a | 44.0 ^{fgh} | 51.0 ^a | 0.81 | <0.001 | <0.001 |
| RDP, % of total N | 56.8 ^{ab} | 52.4 ^f | 54.6 ^{cde} | 52.7 ^{ef} | 49.9 ^{gh} | 53.4 ^{def} | 55.8 ^{abc} | 54.9 ^{bcd} | 51.9 ^{fg} | 57.5 ^a | 56.0 ^{abc} | 49.0 ^h | 0.81 | <0.001 | <0.001 |
| OPA _F results ⁵ | | | | | | | | | | | | | | | |
| FD _{0F} , % of total N | 2.22 ^a | 2.12 ^{cd} | 2.16 ^{bc} | 2.11 ^d | 2.10 ^d | 2.17 ^b | 2.19 ^{ab} | 2.18 ^b | 2.17 ^b | 2.13 ^{cd} | 2.10 ^d | 2.06 ^c | 0.043 | <0.001 | 0.607 |
| θ, per h | 0.130 ^a | 0.096 ^c | 0.111 ^{cd} | 0.097 ^e | 0.086 ^f | 0.104 ^{de} | 0.115 ^{bc} | 0.117 ^{bc} | 0.102 ^e | 0.123 ^{ab} | 0.113 ^c | 0.086 ^f | 0.0036 | <0.001 | <0.001 |
| Adjusted θ, per h | 0.131 ^a | 0.097 ^c | 0.110 ^b | 0.098 ^c | 0.086 ^d | 0.102 ^e | 0.116 ^b | 0.117 ^b | 0.101 ^c | 0.125 ^a | 0.113 ^b | 0.086 ^d | 0.0028 | <0.001 | <0.001 |
| k _d , per h | 0.120 ^a | 0.086 ^c | 0.099 ^b | 0.088 ^c | 0.075 ^d | 0.091 ^c | 0.105 ^b | 0.106 ^b | 0.090 ^c | 0.115 ^a | 0.103 ^b | 0.076 ^d | 0.0028 | <0.001 | <0.001 |
| RUP, % of total N | 41.3 ^{cd} | 45.3 ^b | 42.9 ^c | 45.5 ^b | 48.2 ^a | 45.8 ^b | 42.1 ^{cd} | 42.5 ^{cd} | 47.0 ^{ab} | 41.0 ^d | 43.0 ^c | 48.5 ^a | 0.69 | <0.001 | <0.001 |
| RDP, % of total N | 58.7 ^{ab} | 54.7 ^b | 57.1 ^b | 54.5 ^c | 51.8 ^d | 54.2 ^c | 57.9 ^{ab} | 57.5 ^{ab} | 53.0 ^{cd} | 59.0 ^a | 57.0 ^b | 51.5 ^d | 0.69 | <0.001 | <0.001 |

^{a-h}LSM with different superscript letters among plants are different ($P < 0.05$).

¹Nitrogen composition data repeated from Table 2.

²FD₀ = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$, where OPA = α -phthalaldehyde.

³Soluble protein N = Soluble N - FD₀.

⁴Degradation determined from net release of ammonia plus OPA absorbance (A_{340}), which includes free AA plus oligopeptides.

⁵FD_{0A} = fraction total N present as degraded N [ammonia and free AA plus oligopeptides (A_{340})] at $t = 0$.

⁶Adjusted θ computed by dividing θ observed for individual proteins in each incubation by the ratio: mean incubation θ /overall mean θ .

⁷Degradation rate (k_d) = adjusted $\theta - (FD_0/100)/\text{incubation time (2 h)}$.

⁸RUP = insoluble N $\times 0.06/(0.06 + k_d)$ + soluble protein N $\times 0.16/(0.16 + k_d)$; RDP = 100 - RUP.

⁹Degradation determined from net release of ammonia plus OPA fluorescence, which includes only free AA.

¹⁰FD_{0F} = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at $t = 0$.

Table 7. Estimates of proportions of total variance (%) contributed by sources of variation in statistical model¹

| Item | Source of variation | | | | | | |
|---|---------------------|----------------|------|-------|--------------|-------------|-------|
| | Sample replicate | Incubation run | Year | Plant | Year × plant | Model total | Error |
| Composition ² | | | | | | | |
| Insoluble N | 0.04 | 0.05 | 4.1 | 57.6 | 13.8 | 75.7 | 24.3 |
| Soluble N | 0.04 | 0.05 | 4.1 | 57.6 | 13.8 | 75.7 | 24.3 |
| Soluble protein N ³ | 0.08 | 0.05 | 4.1 | 57.3 | 13.9 | 75.5 | 24.5 |
| OPA _A results ⁴ | | | | | | | |
| FD _{0A} , ⁵ % of total N | 16.9 | 0.12 | 4.7 | 48.2 | 12.6 | 82.5 | 17.5 |
| θ, per h | 0.03 | 8.6 | 32.8 | 19.5 | 9.7 | 70.6 | 29.4 |
| Adjusted θ, ⁶ per h | 0.02 | 0.26 | 0.13 | 34.7 | 16.7 | 51.8 | 48.2 |
| k _d , ⁷ per h | 0.12 | 0.25 | 0.18 | 31.7 | 16.6 | 48.9 | 51.1 |
| RUP, ⁸ % of total N | 0.02 | 0.35 | 0.99 | 20.8 | 19.8 | 41.9 | 58.1 |
| RDP, ⁸ % of total N | 0.02 | 0.35 | 0.99 | 20.8 | 19.8 | 41.9 | 58.1 |
| OPA _F results ⁹ | | | | | | | |
| FD _{0F} , ¹⁰ % of total N | 40.1 | 0.55 | 1.1 | 29.0 | 7.3 | 78.1 | 21.9 |
| θ, per h | 0.05 | 9.5 | 29.8 | 21.8 | 8.5 | 69.6 | 30.4 |
| Adjusted θ, per h | 0.03 | 0.22 | 0.08 | 38.3 | 13.7 | 52.3 | 47.7 |
| k _d , ⁷ per h | 0.06 | 0.21 | 0.07 | 37.9 | 13.7 | 51.9 | 48.1 |
| RUP, ⁸ % of total N | 0.02 | 0.30 | 0.70 | 24.9 | 15.3 | 41.2 | 58.8 |
| RDP, ⁸ % of total N | 0.02 | 0.30 | 0.70 | 24.9 | 15.3 | 41.2 | 58.8 |

¹Proportions of total variance computed by dividing type-3 sums of squares for each source of variation by the total type-3 sums of squares determined in an overall SAS GLM analysis of these items.

²Nitrogen composition data repeated from Table 2.

³Soluble protein N = soluble N – FD₀.

⁴Degradation determined from net release of ammonia plus *o*-phthalaldehyde (OPA) absorbance (A₃₄₀), which includes free AA plus oligopeptides.

⁵FD_{0A} = fraction total N present as degraded N [ammonia and free AA plus oligopeptides (A₃₄₀)] at *t* = 0.

⁶Adjusted θ computed by dividing θ observed for individual proteins in each incubation by the ratio: mean incubation θ/overall mean θ.

⁷Degradation rate (k_d) = adjusted θ – (FD₀/100)/incubation time (2 h).

⁸RUP = insoluble N × 0.06/(0.06 + k_d) + soluble protein N × 0.16/(0.16 + k_d); RDP = 100 – RUP.

⁹Degradation determined from net release of ammonia plus OPA fluorescence, which includes only free AA.

¹⁰FD_{0F} = fraction total N present as degraded N [ammonia and free AA (OPA fluorescence)] at *t* = 0.

ruminal escape with minimal loss of intestinal digestibility. Moreover, greater RUP appears to be supplied by CM produced by some crushing plants than by other CM crushing plants.

CONCLUSIONS

Effects of production year and processing plant on chemical composition and ruminal *in vitro* protein degradation were assessed using a large set of CM samples collected from 12 Canadian crushing plants over 4 yr. Overall mean CM protein degradation rate was 0.093/h and RUP was 45%. Estimates in the same incubations for SSBM and ESBM were, respectively, 0.177 and 0.055/h for degradation rate and 33 and 56% for RUP. Small effects of year were present for CM chemical composition, including N solubility, but differences were partly due to changes in analytical methods over time. However, estimates of CM degradation rate and RUP were not different among processing years. Consistent differences in CM chemical composition, degradation rate, and RUP were observed among processing plants.

Regression analysis indicated that soluble N content explained 40 to 42% of the variation in ruminal degradation rate of CM protein. These results showed important differences in RUP content and thus MP value of CM produced by different canola crushing plants.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the of the barn crew at the University of Wisconsin-Madison Dairy Cattle Center and the US Dairy Forage Research Center Farm (Prairie du Sac, WI) for care and feeding of the rumen donor cows; Wendy Radloff, Mary Becker, Eduardo Marostegan de Paula, and Ian Kraus for assisting with sampling and laboratory analyses; Peter Crump of the University of Wisconsin-Madison for assisting with statistical analyses; and the Canola Council of Canada for partial funding of this project.

REFERENCES

Acharya, I. P., D. J. Schingoethe, K. F. Kalscheur, and D. P. Casper. 2015. Response of lactating dairy cows to dietary protein from

- canola meal or distillers' grains on dry matter intake, milk production, milk composition, and amino acid status. *Can. J. Anim. Sci.* 95:267–279.
- AOAC. 1990. Official Methods of Analysis. 15th ed. Assoc. Off. Anal. Chem., Arlington, VA.
- Block, R. J., and K. W. Weiss. 1956. *The Amino Acid Handbook*. Thomas, Springfield, IL.
- Borucki Castro, S. I., L. E. Phillip, H. Lapierre, P. W. Jardon, and R. Berthiaume. 2007. Ruminal degradability and intestinal digestibility of protein and amino acids in treated soybean meal products. *J. Dairy Sci.* 90:810–822.
- Boucher, S. E., C. Pedersen, H. H. Stein, and C. G. Schwab. 2009. Evaluation of the furosine and homoarginine methods for determining reactive lysine in rumen-undegraded protein. *J. Dairy Sci.* 92:3951–3958.
- Brennan, R. F. 2016. Effect of time of application and amount of nitrogen fertilizer on seed yield and concentration of oil in canola (*Brassica napus*). *J. Plant Nutr.* 39:337–347.
- Brito, A. F., and G. A. Broderick. 2007. Effects of feeding different protein supplements on milk production and nutrient utilization in dairy cows. *J. Dairy Sci.* 90:1816–1827.
- Brito, A. F., G. A. Broderick, and S. M. Reynal. 2007. Effects of different protein supplements on omasal nutrient flow and microbial protein synthesis in lactating dairy cows. *J. Dairy Sci.* 90:1828–1841.
- Broderick, G. A. 1987. Determination of protein degradation rates using a rumen in vitro system containing inhibitors of microbial nitrogen metabolism. *Br. J. Nutr.* 58:463–475.
- Broderick, G. A., and M. K. Clayton. 1992. Rumen protein degradation rates estimated by nonlinear regression analysis of Michaelis-Menten in vitro data. *Br. J. Nutr.* 67:27–42.
- Broderick, G. A., A. P. Faciola, and L. E. Armentano. 2015. Replacing dietary soybean meal with canola meal improves production and efficiency of lactating dairy cows. *J. Dairy Sci.* 98:5672–5687.
- Broderick, G. A., D. B. Ricker, and L. S. Driver. 1990. Expeller soybean meal and corn by-products versus solvent soybean meal for lactating dairy cows. *J. Dairy Sci.* 73:453–462.
- Broderick, G. A., P. Uden, M. L. Murphy, and A. Lapins. 2004. Sources of variation in rates of in vitro ruminal protein degradation. *J. Dairy Sci.* 87:1345–1359.
- Broderick, G. A., R. J. Wallace, and N. McKain. 1988. Uptake of small neutral peptides by mixed rumen microorganisms in vitro. *J. Sci. Food Agric.* 42:109–118.
- Chen, G., C. J. Sniffen, and J. B. Russell. 1987. Concentration and estimated flow of peptides from the rumen of dairy cattle: Effects of protein quantity, protein solubility, and feeding frequency. *J. Dairy Sci.* 70:983–992.
- Choi, C. W., A. Vanhatalo, and P. Huhtanen. 2003. Effects of type of grass silage and level of concentrate on the flow of soluble non-ammonia nitrogen entering the omasum of dairy cows. *J. Anim. Feed Sci.* 12:3–22.
- Classen, H. L., R. W. Newkirk, and D. D. Maenz. 2004. Effects of conventional and novel processing on the feed value of canola meal for poultry. Pages 1–16 in *Proc. Aust. Poultry Sci. Symp.* The Poultry Research Foundation, University of Sydney, Sydney, NSW, Australia.
- Colombini, S., G. A. Broderick, and M. K. Clayton. 2011. Effect of quantifying peptide release on ruminal protein degradation determined using the inhibitor in vitro system. *J. Dairy Sci.* 94:1967–1977.
- Craig, W. M., and G. A. Broderick. 1981. Effect of heat treatment on true digestibility in the rat, in vitro proteolysis and available lysine content of cottonseed meal protein. *J. Anim. Sci.* 52:292–301.
- Faldet, M. A., and L. D. Satter. 1991. Feeding heat-treated full fat soybeans to cows in early lactation. *J. Dairy Sci.* 74:3047–3054.
- Faldet, M. A., L. D. Satter, and G. A. Broderick. 1992. Determining optimal heat treatment of soybeans by measuring available lysine chemically and biologically with rats to maximize protein utilization by ruminants. *J. Nutr.* 122:151–160.
- Friedman, M. 1996. Food browning and its prevention: An overview. *J. Agric. Food Chem.* 44:631–653.
- Hickling, D. 2008. Maximized utilization of canola co-products in livestock industry. Pages 3–14 in *Proc. 29th Western Nutrition Conference*, Edmonton, AB, Canada. University of Alberta, Edmonton, AB, Canada.
- Hintz, R. W., D. R. Mertens, and K. A. Albrecht. 1996. Effects of sodium sulfite on recovery and composition of detergent fiber and lignin. *J. AOAC Int.* 79:16–22.
- Huhtanen, P., M. Hetta, and C. Swensson. 2011. Evaluation of canola meal as a protein supplement for dairy cows: A review and a meta-analysis. *Can. J. Anim. Sci.* 91:529–543.
- Licitra, G., T. M. Hernandez, and P. J. Van Soest. 1996. Standardization of procedures for nitrogen fractionation of ruminant feeds. *Anim. Feed Sci. Technol.* 57:347–358.
- Mahler, H. R., and E. H. Cordes. 1966. *Biological Chemistry*. Harper and Row, New York, NY.
- Martineau, R., D. R. Ouellet, and H. Lapierre. 2013. Feeding canola meal to dairy cows: A meta-analysis on lactational responses. *J. Dairy Sci.* 96:1701–1714.
- Maxin, G., D. R. Ouellet, and H. Lapierre. 2013a. Ruminal degradability of dry matter, crude protein, and amino acids in soybean meal, canola meal, corn, and wheat dried distillers grains. *J. Dairy Sci.* 96:5151–5160.
- Maxin, G., D. R. Ouellet, and H. Lapierre. 2013b. Effect of substitution of soybean meal by canola meal or distillers grains in dairy rations on amino acid and glucose availability. *J. Dairy Sci.* 96:7806–7817.
- McCartney, C. A., R. Scarth, P. B. E. McVetty, and J. K. Daun. 2004. Genotypic and environmental effects on saturated fatty acid concentration of canola grown in Manitoba. *Can. J. Plant Sci.* 84:749–756.
- McDougall, E. I. 1948. Studies on ruminant saliva. I. The composition and output of sheep's saliva. *Biochem. J.* 43:99–109.
- Moshtaghi Nia, S. A., and J. R. Ingalls. 1995. Influence of moist heat treatment on ruminal and intestinal disappearance of amino acids from canola meal. *J. Dairy Sci.* 78:1552–1560.
- Mutsvangwa, T., D. Kiran, and S. Abeysekara. 2016. Effects of feeding canola meal or wheat dried distillers grains with solubles as a major protein source in low- or high-crude protein diets on ruminal fermentation, omasal flow, and production in cows. *J. Dairy Sci.* 99:1216–1227.
- Newkirk, R. 2009. *Canola Meal: Feed Industry Guide*. 4th ed. Canadian International Grains Institute, Winnipeg, MB, Canada.
- Newkirk, R. W., H. L. Classen, T. A. Scott, and M. J. Edney. 2003. The digestibility and content of amino acids in toasted and non-toasted canola meals. *Can. J. Anim. Sci.* 83:131–139.
- NRC. 2001. *Nutrient Requirements of Dairy Cattle*. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- Pritchard, F. M., H. A. Eagles, R. M. Norton, P. A. Salisbury, and M. Nicolas. 2000. Environmental effects on seed composition of Victorian canola. *Aust. J. Exp. Agric.* 40:679–685.
- Roth, M. 1971. Fluorescence reaction of amino acids. *Anal. Chem.* 43:880–882.
- SAS Institute. 2013. *SAS® 9.4-Guide to Software Updates*. SAS Inst. Inc., Cary, NC.
- Segal, I. 1976. *Biochemical Calculations*. 2nd ed. John Wiley & Sons, New York, NY.
- Swanepoel, N., P. H. Robinson, and L. J. Erasmus. 2014. Determining the optimal ratio of canola meal and high protein dried distillers grain protein in diets of high producing Holstein dairy cows. *Anim. Feed Sci. Technol.* 189:41–53.
- Thacker, P. A., and R. W. Newkirk. 2005. Performance of growing-finishing pigs fed barley-based diets containing toasted or non-toasted canola meal. *Can. J. Anim. Sci.* 85:53–59.
- Van Soest, P. J., J. B. Robertson, and B. A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74:3583–3597.
- Waldo, D. R., L. W. Smith, and E. L. Cox. 1972. Model of cellulose disappearance from the rumen. *J. Dairy Sci.* 55:125–129.