

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317750447>

Physical activity, depression and quality of life in aging process

Article · January 2017

CITATIONS

11

READS

351

4 authors, including:



Saniye Aydoğan Arslan

Kirikkale University Faculty of Health Science, Turkey

37 PUBLICATIONS 78 CITATIONS

[SEE PROFILE](#)



Tezel Yıldırım Şahan

University of Health Science Turkey

23 PUBLICATIONS 55 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Physiotherapy [View project](#)



Validity and Reliability of The 3-Meter Backward Walk Test in Individuals with Stroke [View project](#)

Physical activity, depression and quality of life in aging process.

Meral Sertel*, Saniye Aydoğan Arslan, Fatih Kurtoğlu, Tezel Şahan Yıldırım

Department of Physical Therapy and Rehabilitation, Faculty of Health Sciences, Kırıkkale University, Kırıkkale, Turkey

Abstract

Background: In aging process was to show to change physical activity, depression and quality of life values of individuals aged 18-85.

Methods: Four hundred and fifty individuals aged 18-85 who lived in Kırıkkale were included in our study after being determined with the Simple Randomization Sample Method. Three hundred and forty one individuals who were suitable for our inclusion and exclusion criteria completed the study. The individuals were divided into 3 groups: 142 adults (18-45 aged), 137 middle-aged (45-64 aged) and 62 elderly (>65aged). Quality of life values were measured with Nottingham Health Profile (NHP), physical activity levels of the individuals were measured with 24 h Physical Activity Scale (PAS), depression status of individuals were determined with Beck Depression Scale (BDS).

Results: Our study showed that statistically meaningful significance was found when we compared the groups with quality of life and depression. Statistically meaningful significance was specified on PAS measurement in groups with PAS work and sum of PAS variables ($p < 0.005$). There was no difference between PAS and the other variables in groups ($p > 0.005$).

Conclusion: In summary, our results point that elderly individuals have a sedentary and depressive life style. In all groups, physical activity habits support physiological wellness state and health-related quality of life by strengthening physical functionality. Because of that, studies on strengthening and encouraging physical activity must be planned.

Keywords: Older Adults, physical activity, depression, Quality of life.

Accepted on February 15, 2017

Introduction

In recent years, living with increased quality of life has become an important topic as long living became an important topic. Physical activity is one of the criteria for getting older healthily and decreasing health risk that may occur because of age. In 2003, according to the report of World Health Organization (WHO), sedentary life span caused the death of 2 million individuals [1].

Physical activity affects to get better in terms of public health socially and ensuring physical, social, mental and physiological benefits to men, women and everyone in every age [2]. Regular physical activity prevent chronic diseases that from cardiovascular diseases, heart problems, some cancer types and all mortality causes [2,3]. In addition, regular physical activity has positive effects on blood pressure, lipid and lipoprotein profile, weight control, mental health physiological wellness and supports quality of life [4]. 30-60 min middle intensity physical activity a week is also enough to show positive effects on our health [5,6].

Although the benefits of participating to regular activities are known, it has been found in recent studies that physical activity attendance level is lower than the recommended level [7]. For this reason, in recent years, policies and research on

strengthening and encouraging physical activity has been accelerated parallel to the efforts to improve the life quality by the WHO [8]. When the studies in the literature are examined it is observed that as people get older, the life quality decreases. Women's quality of life levels were generally found to be low in studies conducted with healthy adults. In studies conducted with young people, it has been observed that the life quality between 18-24 years of age is lower in spiritual field when compared with the physical field, and again, it has been reported that they have more risky attitudes in terms of psychological health, perceived health and general health [8]. The aim of our study was to compare the physical activity, depression and quality of life values of individuals from different ages.

Methods

18-80 aged women and men who lived in Kırıkkale were included in our study by simple randomization sample method. Our study was announced heard to the volunteers by announcement. Power and Sample Size Program was used for data power analysis and sampling size. At the end of the analysis, it was found that 95% confidence interval and 90% power would be obtained according to error rate 0.05, and the

number of the individuals for each group was determined as 100, and the total number of the cases was determined as 300.

Inclusion criteria of our study were accepted as being cooperative, agreeing to participate in the study and being between 18-80 years of age. Exclusion criteria were having audio-visual problems, having any orthopedically and neurologically problem, having malignancy and chemotherapy, radiotherapy and pregnant women. As a part of the evaluation, interviews were made face-to-face and confirmation forms were taken from each person. Totally, we reached 450 individuals. 341 individual who was appropriate for the inclusion criteria completed the study (Figure 1). Individuals were separated into three groups as 142 adults (18-45 aged), 137 middle age (45-64 aged), 62 elderly (>65 aged). Participants' demographic data were recorded. Individuals' health related quality of life was evaluated with the Nottingham Health Profile Questionnaire (NHP); physical activity levels were examined with 24-hour Physical Activity Scale (PAS); and the depression status was determined with the Beck Depression Scale (BDS).




Figure 1. Constitution of our research groups.

Nottingham Health Profile (NHP) is a questionnaire which evaluates health problems of different parts of life quality. It has two parts. The first part has 38 questions and six sub-categories on physical activity, pain, sleep, social isolation, emotional reactions and energy levels. Individuals are asked to answer the questions by saying "YES" or "NO". For each part, the points vary between 0 and 100. Higher points show more health-related problems. The second part is about factors which do not change with treatment. There is Turkish version of it [9]. 24-Hour Physical Activity Scale (PAS) consists of four parts, which are, job, transport, home and sport activity. The amount of the exercise is explained as being equal to the kilocalorie spent in a minute. But it is preferred that Metabolic Equivalent (MET) is used while calculating the questionnaire points. One Metabolic Equivalent is defined as the energy consumed in a minute while resting and one MET is equal to 1 kcal/kg-hour. While calculating an individual's amount of consumed energy, frequency, density and time of the activity must be known [10].

Beck Depression Scale (BDS) was developed by Beck et al. in 1979. The Turkish version's reliability and validity was performed by Hisli in 1989, and the Cronbach alpha coefficient of this scale was found as 0.80. In an article written on the Turkish version, the cut point was defined as 17. The total point that can be taken from the scale may vary between 0 and 63. 10-16 point is defined as "weak", 17-21 is defined as "middle", 30-63 is defined as "high" depression level [11].

Statistically analysis

The normal distribution of the variables was tested with the Kolmogorov Smirnov Test. The homogeneity of the group variances were defined with the Levene Test. In the tests that met the criteria of pre-conditions of the parametric tests, the T Test and Anova Test were used in independent groups; and the Mann Whitney U Test - Kruskal Wallis Analysis Methods were used in the other conditions. The $P < 0.005$ level was accepted as being statistically meaningful. The defined statistics were given as mean \pm standard deviation (mean \pm SD). The categorical data analyses were evaluated with the Chi-Square test. The analyses of the data were made with the SPSS 17.0 Program (SPSS Inc., Chicago, Illionis, USA).

Results

The ages, body mass index values, BDS, quality of life and physical activity levels of the individuals are shown in Table 1. Statistically significant differences were determined when the groups were compared about the quality of life and depression ($p < 0.005$). While groups were compared with physical activity, differences were found between PAS work and total PAS Scores ($p < 0.005$); however, there was no difference between the other variables of PAS ($p > 0.005$) (Table 1).

Table 1. Individuals socio-demographic, depression, quality of life and physical activity data with different ages.

	Adult (n=142)	Middle Age (n=137)	Elderly (n=62)	p
Age, mean (X \pm SD), year	30.26 \pm 9.60	53.24 \pm 6.43	70.98 \pm 5.82	0.001*
BMI, mean (X \pm SD), kg/m ²	24.56 \pm 4.83	27.64 \pm 4.38	28.72 \pm 5.77	0.001*
Gender, n (%)				0.001*
Women	101(71.1)	49(35.8)	38(61.3)	
Men	41(28.9)	88(64.2)	24(38.07)	
BDS	8.51 \pm 7.15	9.42 \pm 7.46	13.25 \pm 10.07	\pm 0.008*
Nottingham Health Profile				
NHP-pain	5.56 \pm 9.04	17.37 \pm 19.99	39.64 \pm 33.77	\pm 0.001*
NHP-physical function	3.95 \pm 8.53	16.76 \pm 18.72	35.96 \pm 28.16	\pm 0.001*

Physical activity, depression and quality of life in aging process

NHP-fatigue	8.13 ± 12.83	16.83 18.68	± 30.59 25.16	± 0.001*
NHP-sleep	6.32 ± 12.09	17.62 22.86	± 38.92 31.69	± 0.001*
NHP-social isolation	4.77 ± 11.53	12.21 20.26	± 22.12 30.01	± 0.003*
NHP-emotional state	11.17 21.51	± 24.23 28.76	± 44.78 35.68	± 0.001*
Total NHP	39.52 52.33	± 105.16 89.09	± 213.08 142.86	± 0.001*
PAS (24 h)				
Transport	4.75 ± 5.14	4.64 ± 4.66	3.74 ± 3.79	0.399
Work	11.23 ± 9.46	8.95 ± 9.33	5.89 ± 8.80	0.001*
Sport	0.63 ± 1.99	0.79 ± 3.91	0.85 ± 2.58	0.376

Sleep	5.56 ± 2.69	5.71 ± 2.86	5.84 ± 2.53	0.158
Home	10.80 ± 7.29	11.18 ± 8.11	12.20 ± 8.24	0.541
Total	33.98 12.20	± 31.71 16.62	± 28.69 13.23	± 0.022*

*p<0.05; †: Kruskal Wallis Variance Analysis; ††: Chi-Square Test; BDS: Beck Depression Scale; NHP: Nottingham Health Profile; PAS: Physical Activity Scores

In terms of quality of life, the adult group had higher quality of life than the middle and elder group in all the variables of the NHP (p<0.005). When the middle-aged and the elderly groups were compared excluding social isolation parameter of the NHP, statistically significant differences were determined in all parameters of the NHP (pain, physical function, fatigue, sleep, emotional state and total NHP) (p>0.005).

Table 2. Comparison of quality of life and depression Levels On individuals with different ages.

Groups		Adult- Middle Age	Middle Age- Elderly	Elderly- Adults
NHP(P)	z	-5.23	-4.90	-8.85
	p	0.00*	0.00*	0.00*
NHP(PA)	z	-5.95	-4.47	-7.54
	p	0.00*	0.00*	0.00*
NHP(ER)	z	-3.63	-3.69	-6.35
	p	0.00*	0.00*	0.00*
NHP(S)	z	-3.21	-4.75	-7.27
	p	0.00*	0.00*	0.00*
NHP(SI)	z	-2.28	-1.60	-3.20
	p	0.02*	0.10	0.00*
NHP(EL)	z	-2.35	-3.61	-5.26
	p	0.01*	0.00*	0.00*
NHP(Toplam)	z	-7.17	-5.17	-6.16
	p	0.00*	0.00*	0.00*
BDS	z	-1.07	-2.29	-3.07
	p	0.31	0.02*	0.00*

*p<0.05; Mann Whitney-U Test; NHPP: Nottingham Health Profile pain, NHPA: Nottingham Health Profile Physical Activity, NHPER: Nottingham Health Profile Emotional reactions, NHPS: Nottingham Health Profile sleep, NHPSE: Nottingham Health Profile social isolation, NHPES: Nottingham Health Profile energy level, BDS: Beck Depression Scale

When groups were compared in terms of depression status, despite there was no difference between the adults and the middle-aged groups (p>0.005), statistically significant difference was found with the elderly and the middle-aged groups (p<0.005). While middle aged and the elderly groups were compared, the elder participants had higher BDS scores (p<0.005, Table 2). According to PAS total scores, only adults and the elderly groups had statistically significant differences,

which were recorded; and it was observed that adults were more active (p<0.005, Table 3).

Table 3. Comparison of physical activity level on individual's different ages.

Groups	Adult- Middle Age	Middle Elderly	Age- Elderly- Adult
--------	----------------------	-------------------	---------------------------

PAS(Transport)	z	-0.33	-1.31	-1.10
	p	0.73	0.18	0.26
PAS(Work)	z	-1.98	-2.42	-4.03
	p	0.04*	0.01*	0.00*
PAS(Sport)	z	-0.62	-1.40	-0.87
	p	0.53	0.15	0.38
PAS(Sleep)	z	-1.58	-0.43	-1.59
	p	0.11	0.66	0.11
PAS(Home)	z	-0.28	-0.79	-1.15
	p	0.77	0.42	0.24
PAS(Total)	z	-0.58	-1.95	-2.92
	p	0.55	0.05	0.00*

*p<0.05; Mann Whitney-U Test; PAS: 24 h Physical Activity Score

Discussion

There are some cross-sectional and retrospective studies showing that physical activity increases quality of life. Despite many researches investigating the relation between physical activity and depression in general population, there is less research on the effects of physical activity on psychological diseases like depression [12,13]. The aim of our study was to compare physical activity, depression and quality of life values in individuals from different ages.

In our study show that, statistically significant differences were determined between the age and the BMI values in participants. Because of including different ages in the study, it was clear that difference could be found between ages. The BMI values of the elderly and the middle-aged groups were found higher than the adults. According to research, the BMI values of especially middle-aged and the elderly people are increasing [14-17]. The reason of this is the fact that metabolic activity slows down in individuals' body in middle-ages because of having completed growing [16,17]. Decreases in individual's physical movements and basal metabolism speed are formed with aging; this condition causes increases in BMI scores.

In general population, min 40% of people has had a physiological illness at least once in their life spans. In general population, age is a relevant factor for physiopathology. The prevalence of psychiatric disorders and their nature are different in older adults than in middle or young adults [18]. In childhood, attention deficit hyperactivity disorder, behavioral disorder and anxiety disorders are the most prevalent comorbid disorders [19] whereas in adulthood, next to hyperactivity disorder and anxiety disorders, mood disorders are common [20,21]. In various adult studies, older adults with psychiatric disorders have been included but only a few directly compared older adults with younger individuals [21]. Whereas anxiety seemed to reduce from childhood to young adulthood [20], the risk for depression seemed to increase with increasing age [22].

According to the results of our study, the elderly participants were more depressive than the adults and middle-aged participants; furthermore elder participants' quality of life values were lower than the adult participants. Because of aging is a term when individuals have losses in physical appearance, power, role and position and individuals become dependent to their relatives due to physical disorders and loss of functions, approaches of the individuals in this period must be different from the other age groups [23]. Lower quality of life and depressive mood in the elderly people were determined in our study, as well. It was shown in the literature that not only the age factor, but also the gender, socio-economic state (education and work etc.), quality of life, intellectual and cognitive functions are effective on physiological disorders [24]. Lower quality of life and depressive mood in the elderly people are thought to have increased, and the number of the researches on this topic must be increased to investigate the cause of physiological disorders.

Sigmundova et al. showed that older and predominantly sedentary people had higher probabilities of being in the low physical activity level category. Older people and those who are sedentary for more than four hours a day showed increased probabilities of being in the low physical activity level category [25].

Bertrais et al. evaluated leisure time physical activity on 7404 individuals aged between 45 and 68 in their 8-year follow-up study. Physical activity was associated positively with age, gender and education level; but smoking was associated negatively not only in men but also in women [26]. USA data such as those from behavioral risk factor surveillance system, have shown that percentages of subjects recommended that the physical activity levels were higher in younger people (18-29) compared with the other age groups (30-64) [27]. In our results, physical activity levels of the adults were higher than the middle-aged and the elderly groups. It was found that our results supported the researches in the literature. But results of other research for evaluating physical activity habits of different ages present that physical activity is not a life style among children, younger people and the elderly in our country. For this reason, from childhood we must be more active for encouraging regular physical exercise habits [28]. We think that studies on encouraging physical activity in all ages are important for gaining physical activity habits and a healthy life model. As well as these components, there is a need for healthy life programs and precautions with healthy life attitudes in young ages.

Conclusions

In the result of our study, it has been determined that the elderly individuals have a depressive and sedentary lifestyle. Gaining physical activity habits in all ages provides healthy life quality with psychological wellness state and physical functionality. For this reason, studies on strengthening and encouraging physical activity must be planned.

Limitations

Although we included 150 individuals for each group in the beginning of our study, it can be a limitation because of several causes in the elderly group.

Ethical Approval and Consent to Participate

All procedures performed were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. This study was approved by the Kırıkkale University Clinical Research Ethics Committee, Decision Number: 2015/28.

Consent for Publication

Informed consent was obtained from all individual participants included in the study.

References

1. World Health Organization. WHO Global Strategy on Diet, Physical Activity and Health: Eastern Mediterranean Regional Consultation Meeting Report. World Health Organization, Cairo 2003; 1-23.
2. Artikel A. Physical activity and health related quality of life among non-academic staff of a university. *J Sai Kesih Malay* 2015; 13: 69-75.
3. Haskell LW, Lee Mİ, Pate RR. Physical activity and public health: updated recommendation for adults from the american college of sports medicine and the american heart association. *Med. Sci. Sports Exerc* 2007; 39: 1423-1434.
4. Vuori I. Does physical activity enhance health? *Patient Education and Counseling* 1998; 33: 95-103.
5. Nakamura Y, Kikuchi H, Oka K. Stage of change of exercise affects health-related quality of life. *Int J Sport Health Sci* 2006; 4: 67-73.
6. Shibata A, Oka K, Nakamura Y. Recommended level of physical activity and health-related quality of life among Japanese adults. *Heal Qual of Life Outcom* 2007; 5: 64.
7. Vuillemin A, Boini S, Bertrais S. Leisure time physical activity and health related quality of life. *Preven Med* 2005; 41: 562-569.
8. Bize R, Johnson J, Plotnikoff R. Physical activity level and healthrelated quality of life in the general adult population: a systematic review. *Prev Med* 2007; 45: 401-415.
9. Küçükdeveci AA, McKenna SP, Kutlay S. The development and psychometric assessment of the Turkish version of the Nottingham Health Profile. *Int J Rehabil Res* 2000; 23: 31-38.
10. Karaca A. Reliability and validity of three different questionnaires in employees. Hacettepe University Health Sciences Institute, PhD Thesis, in School of Sports Sciences and Technology Program, Ankara. 2004.
11. Hisli N. Beck Depresyon Envanterinin üniversite öğrencileri için geçerliliği güvenilirliği. *Türk Psikoloji Dergisi* 1989; 7: 3-13.
12. Eime R, Harvey J, Payne W. Dose-response of women's health-related quality of life (HRQoL) and life satisfaction to physical activity. *J Phys Act Health* 2014; 11: 330-338.
13. Wolin K, Glynn R, Colditz G. Long-term physical activity patterns and health-related quality of life in US women. *Am J Prev Med* 2007; 32: 490-499.
14. Aktener YA, Dülger İH, Erkayhan EG. Obesity prevalence in reproductive age and postmenopausal women aged between 20-64 years in a semiurban area. *Trakya Ü. Tıp Fak. Dergisi* 2006; 23: 119-126.
15. Koruk İ, Şahin KT. The prevalence of obesity and risk factors in housewives 15-49 age group in region Konya Fazilet Uluişik health center. *Genel Tıp Dergisi* 2005; 15: 147-155.
16. Okyay P, Uçku R. Prevalence and risk factors of obesity of reproductıve age women in an Urban Area of Izmir. *ADÜ Tıp Fak. Dergisi* 2002; 3: 5-12.
17. Doğan N, Toprak D, Demir S. Prevalence of obesity and associated risk factors in Afyonkarahisar-Turkey. *Türkiye Klinikleri J Med Sci* 2011; 31: 122-32.
18. Kessler RC, Berglund P, Demler O. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Arch of Gen Psych* 2005; 62: 593-602.
19. Simonoff E, Pickles A, Charman T. Psychiatric disorders in children with autism spectrum disorders: Prevalence, comorbidity, and associated factors in a population-derived sample. *J Am Acad Child Adolesc Psychiatry* 2008; 47: 921-929.
20. Croen LA, Zerbo O, Qian Y. The health status of adults on the autism spectrum. *Autism. The International Journal of Research and Practice* 2015; 19: 814-823.
21. Roy M, Prox-Vagedes V, Ohlmeier MD. Beyond childhood: Psychiatric comorbidities and social background of adults with asperger syndrome. *Psychiatria Danubina* 2015; 27: 50-59.
22. Ghaziuddin M, Ghaziuddin N, Greden J. Depression in persons with autism: implications for research and clinical care. *J Autism Dev Disord* 2002; 32: 299-306.
23. Aydemir Ç. Türkiye'de yaşlı intiharları. *Kriz Dergisi* 1999; 7: 21- 25.
24. Lever AG, Geurts HM. Psychiatric co-occurring symptoms and disorders in young, middle-aged, and older adults with autism spectrum disorder. *J Autism Dev Disord*.
25. Sigmundová D, Sigmund E, Hamřík Z. Sedentary behaviour and physical activity of randomised sample of czech adults aged 20-64 years: İpaq And Gpaq Studies Between 2002 And 2011. *Cent Eur J Public Health* 2015; 23: 91-96.
26. Bertrais S, Preziosi P, Mennen L. Socio demographic and geographic correlates of meeting current recommendations for physical activity in middle-aged French adults: the Supplémentation en Vitamines et Minéraux Antioxydants (SUVIMAX) Study. *Am J Public Health* 2004; 94: 1560-6.
27. Macera CA, Pratt M. Public health surveillance of physical activity. *Res Q Exerc Sport* 2000; 71: 97-103.

28. Gül Baltacı, Irmak H, Kesici C. Fiziksel aktivite bilgi serisi.
Klasmat Matbacılık, Ankara 2008.

***Correspondence to:**

Meral Sertel

Kırıkkale University,

Faculty of Health Sciences,

Department of Physical Therapy and Rehabilitation,

Kırıkkale,

Turkey