



Miskolc Mathematical Notes
Vol. 14 (2013), No 3, pp. 1021-1030

HU e-ISSN 1787-2413
DOI: 10.18514/MMN.2013.531

On bivariate Meyer-König and Zeller operators

Ali Olgun



ON BIVARIATE MEYER-KÖNIG AND ZELLER OPERATORS

ALI OLGUN

Received 3 May, 2012

Abstract. This work relates to bivariate Meyer-König and Zeller operators, M_n , $n \in \mathbb{N}$ which are not a tensor product setting. We show the monotonicity of the sequence of operators for n under convexity, moreover we study the property of monotonicity in the sense of Li [9]. Finally, we provide an r th order generalization $M_n^{[r]}$ of M_n and also study approximation of $M_n^{[r]}$.

2010 *Mathematics Subject Classification:* 41A25; 41A36

Keywords: multivariate Meyer-König and Zeller operator, convexity, monotonicity, modulus of continuity

1. INTRODUCTION

The Cheney and Sharma modification of the well-known univariate Meyer-König and Zeller operators (MKZ) are defined as

$$M_n^*(f, x) = \sum_{k=0}^{\infty} m_{n,k}(x) f\left(\frac{k}{n+k}\right), \quad n \in \mathbb{N} \quad (1.1)$$

for $f \in C[0, 1)$ and $x \in [0, 1)$, $n \in \mathbb{N}$, where $m_{n,k}(x) = (1-x)^{n+1} \binom{n+k}{k} x^k$ [3].

The monotonic convergence of $\{M_n^*(f; x)\}_{n=1}^{\infty}$ under convexity was investigated in [2] by Cheney and Sharma by means of analytical approach, also studied in [8] by Khan, by means of probabilistic approach. Monotonic convergence of several approximation operators are studied deeply in [6] by Khan, Della-Vecchia and Fassih with probabilistic point of view. Some works related to MKZ operators can be viewed in [1–8, 13] and [10, 11].

We shall use the following standard notation. Let $\mathbf{x} = (x_1, x_2)$, $\mathbf{t} = (t_1, t_2) \in \mathbb{R}^2$, $\mathbf{k} = (k_1, k_2) \in \mathbb{N}_0^2$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbf{x}^{\mathbf{k}} = x_1^{k_1} \cdot x_2^{k_2}$, $\mathbf{k}! = k_1! \cdot k_2!$, $|\mathbf{k}| = k_1 + k_2$, $|\mathbf{x}| =$

$x_1 + x_2$, \mathbf{e}_i denotes the unit vector in \mathbb{R}^2 . Furthermore $\binom{n}{\mathbf{k}} = \frac{n!}{\mathbf{k}!(n-|\mathbf{k}|)!}$, and

$$\sum_{\mathbf{k}=0}^{\infty} = \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty}.$$

Let $S_2 \subset \mathbb{R}^2$ be the open simplex defined by

$$S_2 = \{\mathbf{x} \in \mathbb{R}^2; \mathbf{x}_i \geq 0, i = 1, 2, |\mathbf{x}| < 1\}.$$

For a function $f \in C(S_2)$, bivariate MKZ operators are defined as

$$M_n(f, \mathbf{x}) = (1 - |\mathbf{x}|)^{n+1} \sum_{\mathbf{k}=0}^{\infty} f\left(\frac{\mathbf{k}}{n + |\mathbf{k}|}\right) \binom{n + |\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}}, \quad n \in \mathbb{N}, \quad (1.2)$$

where $C(S_2)$ denotes the space of continuous real valued functions defined on S_2 . It is obvious that M_n , $n \in \mathbb{N}$, are not a tensor product extension of the univariate MKZ operators M_n^* given by (1.2).

Now we give the following definitions which we shall use.

Definition 1. A continuous function f is said to be convex in $D \subset \mathbb{R}^m$, if

$$f\left(\sum_{i=1}^r \alpha_i \mathbf{x}_i\right) \geq \sum_{i=1}^r \alpha_i f(\mathbf{x}_i)$$

for every $x_1, x_2, \dots, x_r \in D$ and for every non-negative numbers $\alpha_1, \alpha_2, \dots, \alpha_r$ such that $\alpha_1 + \alpha_2 + \dots + \alpha_r = 1$.

Definition 2. A continuous function f from $D \subseteq \mathbb{R}^2$ into \mathbb{R} is said to be Lipschitz continuous of order μ , $\mu \in (0, 1]$, if there exists a constant $A > 0$ such that for every $(x_1, x_2), (y_1, y_2) \in D$, f satisfies

$$|f(x_1, x_2) - f(y_1, y_2)| \leq A \sum_{i=1}^2 |x_i - y_i|^{\mu},$$

the set of Lipschitz continuous functions is denoted by $Lip_A(\mu, D)$.

In this work, we firstly show the monotonicity of the sequence of bivariate MKZ operators defined by (1.2) under convexity. Secondly we give a kind of monotonicity similar to the property given by Li in [9]. Namely, we show that if $f(\mathbf{x})$ is a non-negative function and $x_i^{-1} f(\mathbf{x})$ ($i = 1, 2$) is non-increasing for x_i on $(0, 1)$, then for each $n \geq 1$, $x_i^{-1} M_n(f; \mathbf{x})$ is also non-increasing for x_i on $(0, 1)$. Moreover we build an r -th order generalization $M_n^{[r]}$ of M_n analogues to Kirov and Popova's construction in [8] and investigate its approximation property.

2. MONOTONICITY FOR THE SEQUENCE OF BIVARIATE MEYER-KÖNIG AND ZELLER OPERATORS

In this section, we study the monotonic convergence of the sequence of bivariate MKZ operator under convexity. Note that monotonic convergence of univariate MKZ operator when f is convex, was first obtained by Cheney and Sharma in [3]. We note here that monotonic convergence of the multivariate Baskakov operator is studied in [2].

Theorem 1. *If f is convex, then $M_n(f; \mathbf{x})$ is strictly monotonically non-decreasing in n , unless f is the linear function (in which case $M_n(f; \mathbf{x}) = M_{n+1}(f; \mathbf{x})$ for all $n \in \mathbb{N}$).*

Proof. We have

$$\begin{aligned}
 M_n(f; \mathbf{x}) - M_{n+1}(f; \mathbf{x}) &= (1 - |\mathbf{x}|)^{n+1} \times \\
 &\times \left\{ \sum_{k_2=1}^{\infty} \left[f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} - f\left(0, \frac{k_2}{n+1+k_2}\right) \binom{n+1+k_2}{k_2} \right] \mathbf{x}^{\mathbf{k}} \right. \\
 &+ \sum_{k_1=1}^{\infty} \left[f\left(\frac{k_1}{n+k_1}, 0\right) \binom{n+k_1}{k_1} - f\left(\frac{k_1}{n+1+k_1}, 0\right) \binom{n+1+k_1}{k_1} \right] \mathbf{x}^{\mathbf{k}} \\
 &+ \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{\mathbf{k}}{n+1+|\mathbf{k}|}\right) \binom{n+1+|\mathbf{k}|}{\mathbf{k}} x_1^{k_1+1} x_2^{k_2} \\
 &\left. + \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{\mathbf{k}}{n+1+|\mathbf{k}|}\right) \binom{n+1+|\mathbf{k}|}{\mathbf{k}} x_1^{k_1} x_2^{k_2+1} \right\}.
 \end{aligned}$$

Therefore the last equation reduces to the following:

$$\begin{aligned}
 M_n(f; \mathbf{x}) - M_{n+1}(f; \mathbf{x}) &= (1 - |\mathbf{x}|)^{n+1} \times \\
 &\times \left\{ \sum_{k_2=1}^{\infty} \left[f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} - f\left(0, \frac{k_2}{n+1+k_2}\right) \binom{n+1+k_2}{k_2} \right] \right. \\
 &+ f\left(0, \frac{k_2-1}{n+k_2}\right) \binom{n+k_2}{k_2-1} x_2^{k_2} + \sum_{k_1=1}^{\infty} \left[f\left(\frac{k_1}{n+k_1}, 0\right) \binom{n+k_1}{k_1} \right. \\
 &- f\left(\frac{k_1}{n+1+k_1}, 0\right) \binom{n+1+k_1}{k_1} + f\left(\frac{k_1-1}{n+k_1}, 0\right) \binom{n+k_1}{k_1-1} \left. \right] x_1^{k_1} \\
 &\left. + \sum_{k_1=1}^{\infty} \sum_{k_2=1}^{\infty} \left[f\left(\frac{k_1-1}{n+|\mathbf{k}|}, \frac{k_2}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}-e_1} \right] \right\}.
 \end{aligned}$$

$$\begin{aligned}
& + f\left(\frac{k_1}{n+|\mathbf{k}|}, \frac{k_2-1}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}-\mathbf{e}_2} \\
& + f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}} - f\left(\frac{\mathbf{k}}{n+1+|\mathbf{k}|}\right) \binom{n+1+|\mathbf{k}|}{\mathbf{k}} \Big] \mathbf{x}^{\mathbf{k}} \Big\}.
\end{aligned}$$

Let I_1 , I_2 and I_3 denote the followings

$$\begin{aligned}
I_1 &= f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}} - f\left(\frac{\mathbf{k}}{n+1+|\mathbf{k}|}\right) \binom{n+1+|\mathbf{k}|}{\mathbf{k}} \\
& + f\left(\frac{\mathbf{k}-\mathbf{e}_1}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}-\mathbf{e}_1} + f\left(\frac{\mathbf{k}-\mathbf{e}_2}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}-\mathbf{e}_2},
\end{aligned}$$

$$\begin{aligned}
I_2 &= f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} - f\left(0, \frac{k_2}{n+1+k_2}\right) \binom{n+1+k_2}{k_2} \\
& + f\left(0, \frac{k_2-1}{n+k_2}\right) \binom{n+k_2}{k_2-1},
\end{aligned}$$

$$\begin{aligned}
I_3 &= f\left(\frac{k_1}{n+k_1}, 0\right) \binom{n+k_1}{k_1} - f\left(\frac{k_1}{n+1+k_1}, 0\right) \binom{n+1+k_1}{k_1} \\
& + f\left(\frac{k_1-1}{n+k_1}, 0\right) \binom{n+k_1}{k_1-1}.
\end{aligned}$$

For I_1 , we take

$$\alpha_1 = \frac{\binom{n+|\mathbf{k}|}{\mathbf{k}}}{\binom{n+1+|\mathbf{k}|}{\mathbf{k}}}, \quad \alpha_2 = \frac{\binom{n+|\mathbf{k}|}{\mathbf{k}-\mathbf{e}_1}}{\binom{n+1+|\mathbf{k}|}{\mathbf{k}}}, \quad \alpha_3 = \frac{\binom{n+|\mathbf{k}|}{\mathbf{k}-\mathbf{e}_2}}{\binom{n+1+|\mathbf{k}|}{\mathbf{k}}}.$$

Clearly α_1 , α_2 and α_3 are non-negative numbers, and $\alpha_1 + \alpha_2 + \alpha_3 = 1$. On the other hand if we set

$$\mathbf{x}_1 = \frac{\mathbf{k}}{n+|\mathbf{k}|}, \quad \mathbf{x}_2 = \frac{\mathbf{k}-\mathbf{e}_1}{n+|\mathbf{k}|}, \quad \mathbf{x}_3 = \frac{\mathbf{k}-\mathbf{e}_2}{n+|\mathbf{k}|},$$

then it follows that

$$\alpha_1 \mathbf{x}_1 + \alpha_2 \mathbf{x}_2 + \alpha_3 \mathbf{x}_3 = \frac{\mathbf{k}}{n+1+|\mathbf{k}|}.$$

Hence, from the convexity of f we obtain that $I_1 \leq 0$. For I_2 , we set

$$\alpha_1 = \frac{\binom{n+k_2}{k_2}}{\binom{n+1+k_2}{k_2}}, \quad \alpha_2 = \frac{\binom{n+k_2}{k_2-1}}{\binom{n+1+k_2}{k_2}}$$

and

$$\mathbf{y}_1 = \left(0, \frac{k_2}{n+k_2}\right), \quad \mathbf{y}_2 = \left(0, \frac{k_2-1}{n+k_2}\right),$$

then, clearly, $\alpha_1, \alpha_2 \geq 0$, and $\alpha_1 + \alpha_2 = 1$. Moreover we have

$$\alpha_1 \mathbf{y}_1 + \alpha_2 \mathbf{y}_2 = \left(0, \frac{k_2}{n+k_2}\right).$$

Thus, from the convexity of f we obtain that $I_2 \leq 0$. Similarly we deduce that $I_3 \leq 0$. So we have proved $M_n(f; \mathbf{x}) \leq M_{n+1}(f; \mathbf{x})$ for all $n \in \mathbb{N}$. Using the similar arguments given in [2], the last part of the proof is given as follows: The equality of $M_n(f; \mathbf{x})$ and $M_{n+1}(f; \mathbf{x})$ can be satisfied only if $I_1 = 0$ for all $k_1, k_2 \in \mathbb{N}$ and $I_2 = I_3 = 0$ for all $k_1, k_2 \in \mathbb{N}_0$. But, if f is convex, then $I_1 = 0$ implies that its graph is a plane in triangle. \square

In the following, we show that the bivariate MKZ operators can retain a certain monotony, that is

Theorem 2. *Let f be defined on S_2 . If $f(\mathbf{x})$ is a non-negative function such that $\frac{f(\mathbf{x})}{x_i}$ ($i = 1, 2$) is non-increasing for x_i on $(0, 1)$, then for each $n \geq 1$, $\frac{M_n(f; \mathbf{x})}{x_i}$, is also non-increasing for x_i on $(0, 1)$.*

Proof. Straightforward computation gives that for $n \geq 1$ we have, for example with respect to x_1 ,

$$\begin{aligned} & \frac{\partial}{\partial x_1} \left(\frac{M_n(f; \mathbf{x})}{x_1} \right) = \\ & = \frac{\partial}{\partial x_1} \left\{ \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \frac{(1-x_1-x_2)^{n+1}}{x_1} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} x_1^{k_1} x_2^{k_2} \right\} \\ & = \frac{\partial}{\partial x_1} \left\{ \sum_{k_2=0}^{\infty} \frac{(1-x_1-x_2)^{n+1}}{x_1} f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} \right. \\ & \times (1-x_1-x_2)^{n+1} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} x_1^{k_1-1} x_2^{k_2} \left. \right\} \\ & = \sum_{k_2=0}^{\infty} \frac{\partial}{\partial x_1} \left(\frac{(1-x_1-x_2)^{n+1}}{x_1} \right) f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} \end{aligned}$$

$$\begin{aligned}
& \times f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} \frac{\partial}{\partial x_1} \left[(1-x_1-x_2)^{n+1} x_1^{k_1-1} x_2^{k_2} \right] \\
& = \sum_{k_2=0}^{\infty} \left(\frac{-(n+1)(1-x_1-x_2)^n x_1 - (1-x_1-x_2)^{n+1}}{x_1^2} \right) f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} \\
& + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} x_2^{k_2} \\
& \times \left[-(n+1)(1-x_1-x_2)^n x_1^{k_1-1} + (1-x_1-x_2)^{n+1} (k_1-1) x_1^{k_1-2} \right] \\
& = \sum_{k_2=0}^{\infty} \frac{(1-x_1-x_2)^{n+1}}{x_1^2} \left(\frac{-(n+1)x_1}{1-x_1-x_2} - 1 \right) f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} \\
& + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} (1-x_1-x_2)^n x_2^{k_2} \\
& \times \left[-n x_1^{k_1-1} - x_1^{k_1-1} + (k_1-1) x_1^{k_1-2} - (k_1-1) x_1^{k_1-1} - (k_1-1) x_1^{k_1-2} x_2 \right] \\
& \times x_1^{k_1-2} (1-x_2) (1-x_1-x_2)^n x_2^{k_2} \\
& = \sum_{k_2=0}^{\infty} \frac{(1-x_1-x_2)^{n+1}}{x_1^2} \left(-\frac{(1+nx_1-x_2)}{1-x_1-x_2} \right) f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} \\
& + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} (1-x_1-x_2)^n x_2^{k_2} \\
& \times \left[-(n+k_1) x_1^{k_1-1} + (k_1-1) (1-x_2) x_1^{k_1-2} \right] \\
& = \sum_{k_2=0}^{\infty} \frac{(1-x_1-x_2)^n}{x_1^2} (-(1+nx_1-x_2)) f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} x_2^{k_2} \\
& + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} x_1^{k_1-1} (-n-k_1) x_2^{k_2} \\
& \times (1-x_1-x_2)^n + \sum_{k_1=2}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} \\
& = \sum_{k_2=0}^{\infty} \frac{-(1+nx_1-x_2)}{x_1^2} f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} (1-x_1-x_2)^n x_2^{k_2} \\
& - \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} x_1^{k_1-1} (n+k_1) x_2^{k_2}
\end{aligned}$$

$$\begin{aligned} &\times (1-x_1-x_2)^n + \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1+1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+1+k_2}{k_1+1+k_2} \\ &\times (1-x_2)x_1^{k_1-1}k_1(1-x_1-x_2)^n x_2^{k_2} \\ &\leq - \sum_{k_2=0}^{\infty} \frac{(1+nx_1-x_2)}{x_1^2} f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} (1-x_1-x_2)^n x_2^{k_2} \\ &+ \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1+1}{n+k_1+k_2+1}, \frac{k_2}{n+k_1+k_2+1}\right) \binom{n+k_1+k_2}{k_1+k_2} \frac{n+k_1+k_2+1}{k_1+k_2+1} \frac{k_1}{x_1} \\ &\times (1-x_2)(1-x_1-x_2)^n x_1^{k_1} x_2^{k_2} - \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \binom{n+k_1+k_2}{k_1+k_2} \\ &\times \frac{(n+k_1)}{k_1} \frac{k_1}{x_1} (1-x_1-x_2)^n x_1^{k_1} x_2^{k_2} \end{aligned}$$

Since for $k_1, k_2 \geq 0$

$$\frac{n+k_1+k_2+1}{k_1+k_2+1} < \frac{n+k_1+k_2+1}{k_1+1}, \quad -\frac{n+k_1+k_2}{k_1} < -\frac{n+k_1}{k_1}$$

and

$$1+nx_1-x_2 \geq 0, \quad 1-x_2 \leq 1, \quad \text{for } x_1, x_2 \in S_2, \quad n \in \mathbb{N},$$

so we have for $n \geq 1$

$$\begin{aligned} &\frac{\partial}{\partial x_1} \left(\frac{M_n(f; \mathbf{x})}{x_1} \right) \leq \\ &\leq - \sum_{k_2=0}^{\infty} \frac{(1+nx_1-x_2)}{x_1^2} f\left(0, \frac{k_2}{n+k_2}\right) \binom{n+k_2}{k_2} (1-x_1-x_2)^n x_2^{k_2} \\ &+ \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} \left\{ f\left(\frac{k_1+1}{n+k_1+k_2+1}, \frac{k_2}{n+k_1+k_2+1}\right) \frac{n+k_1+k_2+1}{k_1+k_2+1} \right. \\ &\quad \left. - f\left(\frac{k_1}{n+k_1+k_2}, \frac{k_2}{n+k_1+k_2}\right) \frac{(n+k_1+k_2)}{k_1} \right\} \\ &\times \binom{n+k_1+k_2}{k_1+k_2} \frac{k_1}{x_1} (1-x_1-x_2)^n x_1^{k_1} x_2^{k_2}, \end{aligned}$$

is non-positive, since f is a non-negative function, such that $\frac{f(\mathbf{x})}{x_i}$ ($i = 1, 2$) is non-increasing for x_i on $(0, 1)$. Similar calculations can be obtained for x_2 . □

3. A GENERALIZATION

This section provides an r -th order generalization of the bivariate MKZ operators in the sense of Kirov and Popova's construction [8].

Let $C^r(S_2)$, $r \in \mathbb{N}_0$, denote the space of all functions f defined on S_2 and having all continuous partial derivatives up to order r . By $M_n^{[r]}$, we denote the following generalization of M_n . For $\mathbf{x}, \mathbf{t} \in S_2$,

$$M_n^{[r]}(f; \mathbf{x}) = M_n \left(P_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) \right), \quad (3.1)$$

where

$$\begin{aligned} \Delta \mathbf{x} &:= (\Delta x_1, \Delta x_2) = \mathbf{x} - \frac{\mathbf{k}}{n+|\mathbf{k}|} = \left(x_1 - \frac{k_1}{n+|\mathbf{k}|}, x_2 - \frac{k_2}{n+|\mathbf{k}|} \right), \\ \nabla &= \left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2} \right), \\ (\Delta \mathbf{x} \cdot \nabla)^r &:= \sum_{i+j=r} \binom{r}{j} (\Delta x_1)^i (\Delta x_2)^j \frac{\partial^r}{\partial x_1^i \partial x_2^j}, \end{aligned} \quad (3.2)$$

$\binom{r}{j}$ are binomial coefficients, and

$$\begin{aligned} P_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) &= f \left(\frac{\mathbf{k}}{n+|\mathbf{k}|} \right) + (\Delta \mathbf{x} \cdot \nabla) f \left(\frac{\mathbf{k}}{n+|\mathbf{k}|} \right) \\ &+ \frac{(\Delta \mathbf{x} \cdot \nabla)^2}{2!} f \left(\frac{\mathbf{k}}{n+|\mathbf{k}|} \right) + \dots + \frac{(\Delta \mathbf{x} \cdot \nabla)^r}{r!} f \left(\frac{\mathbf{k}}{n+|\mathbf{k}|} \right), \end{aligned} \quad (3.3)$$

the Taylor polynomial for f at $\frac{\mathbf{k}}{n+|\mathbf{k}|} \in S_2$.

Now we state the following pointwise estimate for $M_n^{[r]}$

Theorem 3. *If $f \in C^r(S_2)$ and $\frac{\partial^r f}{\partial x_1^i \partial x_2^j} \in Lip_A(\gamma)$, $i + j = r$, then we have*

$$\left| M_n^{[r]}(f; \mathbf{x}) - f(\mathbf{x}) \right| \leq \frac{2A}{(r-1)!} \frac{\gamma}{\gamma+r} B(\gamma, r) M_n(g; \mathbf{x}) \quad (3.4)$$

for $x \in S_2$, where $g(\mathbf{s}) = |\mathbf{x} - \mathbf{s}|^{r+\gamma}$, $B(\gamma, r)$ is the familiar beta function, $r, n \in \mathbb{N}_0$, $0 < \gamma \leq 1$ and $A > 0$.

Proof. From (3.1) and (3.3) we have

$$f(\mathbf{x}) - M_n^{[r]}(f; \mathbf{x}) = \sum_{\mathbf{k}=0}^{\infty} R_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) \binom{n+|\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1-|\mathbf{x}|)^{n+1}, \quad (3.5)$$

where

$$R_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) := f(\mathbf{x}) - \sum_{h=0}^r \frac{(\Delta \mathbf{x} \cdot \nabla)^h}{h!} f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|}\right) \binom{n+|\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1-|\mathbf{x}|)^{n+1}. \tag{3.6}$$

(3.6) can be given by

$$R_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) = \frac{1}{(r-1)!} \int_0^1 (\Delta \mathbf{x} \cdot \nabla)^r \times \left[f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|} + t \Delta \mathbf{x}\right) - f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|}\right) \right] (1-t)^{r-1} dt. \tag{3.7}$$

Using (3.2),(3.7) results in

$$R_{r, \frac{\mathbf{k}}{n+|\mathbf{k}|}}(\Delta \mathbf{x}, f) = \frac{1}{(r-1)!} \int_0^1 \sum_{i+j=r} \binom{r}{j} (\Delta x_1)^i (\Delta x_2)^j \frac{\partial^r}{\partial x_1^i \partial x_2^j} \times \left[f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|} + t \Delta \mathbf{x}\right) - f\left(\frac{\mathbf{k}}{n+|\mathbf{k}|}\right) \right] (1-t)^{r-1} dt. \tag{3.8}$$

Substituting (3.8) into (3.5) and taking into account that $\frac{\partial^r f}{\partial x_1^i \partial x_2^j} \in Lip_A(\gamma)$ we arrive at the following.

$$\begin{aligned} \left| f(\mathbf{x}) - M_n^{[r]}(f; \mathbf{x}) \right| &\leq \frac{A}{(r-1)!} \sum_{\mathbf{k}=0}^{\infty} (|\Delta x_1| + |\Delta x_2|)^r (|\Delta x_1|^\gamma + |\Delta x_2|^\gamma) \\ &\times \binom{n+|\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1-|\mathbf{x}|)^{n+1} \int_0^1 t^\gamma (1-t)^{r-1} dt \\ &\leq \frac{2A}{(r-1)!} \sum_{\mathbf{k}=0}^{\infty} (|\Delta x_1| + |\Delta x_2|)^{r+\gamma} \binom{n+|\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1-|\mathbf{x}|)^{n+1} B(\gamma+1, r) \\ &\leq \frac{2A}{(r-1)!} \frac{\gamma}{\gamma+r} B(\gamma, r) \sum_{\mathbf{k}=0}^{\infty} \left| \mathbf{x} - \frac{\mathbf{k}}{n+|\mathbf{k}|} \right|^{r+\gamma} \binom{n+|\mathbf{k}|}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1-|\mathbf{x}|)^{n+1}, \end{aligned}$$

which proves (3.4). □

For the uniform convergence of $M_n^{[r]}(f)$, let us consider the above mentioned function $g(\mathbf{s}) = |\mathbf{x} - \mathbf{s}|^{r+\gamma}$. Obviously $g(\mathbf{x}) = 0$ and g is continuous on S_2 . From multivariate extension of the Bohman-Korovkin theorem (see [12]) we have

$$\|M_n(g)\|_{C(S_2)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore by means of Theorem 3 we arrive at the following result

$$\left\| M_n^{[r]}(f) - f \right\|_{C(S_2)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

REFERENCES

- [1] A. Altın, O. Dođru, and F. Taşdelen, “The generalization of Meyer-König and Zeller operators by generating functions,” *J. Math. Anal. Appl.*, vol. 312, no. 1, pp. 181–194, 2005.
- [2] F. Cao, C. Ding, and Z. Xu, “On multivariate Baskakov operator,” *J. Math. Anal. Appl.*, vol. 307, no. 1, pp. 274–291, 2005.
- [3] E. W. Cheney and A. Sharma, “Bernstein power series,” *Can. J. Math.*, vol. 16, pp. 241–252, 1964.
- [4] O. Dođru and O. Duman, “Statistical approximation of Meyer-König and Zeller operators based on q -integers,” *Publ. Math.*, vol. 68, no. 1-2, pp. 199–214, 2006.
- [5] O. Dođru and V. Gupta, “Korovkin-type approximation properties of bivariate q -Meyer-König and Zeller operators,” *Calcolo*, vol. 43, no. 1, pp. 51–63, 2006.
- [6] M. Khan, B. Della Vecchia, and A. Fassih, “On the monotonicity of positive linear operators,” *J. Approx. Theory*, vol. 92, no. 1, pp. 22–37, art. no. at963 113, 1998.
- [7] R. A. Khan, “Some probabilistic methods in the theory of approximation operators,” *Acta Math. Acad. Sci. Hung.*, vol. 35, pp. 193–203, 1980.
- [8] G. Kirov and L. Popova, “A generalization of the linear positive operators,” *Math. Balk., New Ser.*, vol. 7, no. 2, pp. 149–162, 1993.
- [9] Z. Li, “Bernstein polynomials and modulus of continuity,” *J. Approx. Theory*, vol. 102, no. 1, pp. 171–174, art. no. jath.1999.3374, 2000.
- [10] A.-J. López-Moreno and F.-J. Muñoz-Delgado, “Asymptotic expansion of multivariate conservative linear operators,” *J. Comput. Appl. Math.*, vol. 150, no. 2, pp. 219–251, 2003.
- [11] T. Trif, “An elementary proof of the preservation of Lipschitz constants by the Meyer-König and Zeller operators,” *JIPAM, J. Inequal. Pure Appl. Math.*, vol. 4, no. 5, p. 3, 2003.
- [12] V. I. Volkov, “On the convergence of sequences of linear positive operators in the space of continuous functions of two variables,” *Dokl. Akad. Nauk SSSR*, vol. 115, pp. 17–19, 1957.
- [13] H. Wang, “Properties of convergence for the q -Meyer-König and Zeller operators,” *J. Math. Anal. Appl.*, vol. 335, no. 2, pp. 1360–1373, 2007.

Author’s address

Ali Olgun

Kırıkkale University, Faculty of Art and Science, Department of Mathematics, 71450, Yahşihan, Kırıkkale, Turkey

E-mail address: aliolgun71@gmail.com