

Journal section: Oral Medicine and Pathology
 Publication Types: Research

doi:10.4317/medoral.22520
<http://dx.doi.org/doi:10.4317/medoral.22520>

Investigation of SOSTDC1 gene in non-syndromic patients with supernumerary teeth

Volkan Arikan ¹, Ozge Cumaogullari ², Betul-Memis Ozgul ³, Firdevs-Tulga Oz ⁴

¹ DDS, PhD, Assistant Professor, Department of Pediatric Dentistry, Faculty of Dentistry, Kirikkale University, Kirikkale, Turkey

² PhD, Biotechnology Institute, Ankara University, Ankara, Turkey

³ DDS, PhD, Assistant Professor, Department of Pediatric Dentistry, Faculty of Dentistry, Baskent University, Ankara, Turkey

⁴ DDS, PhD, Professor, Department of Pediatric Dentistry, Faculty of Dentistry, Ankara University, Ankara, Turkey

Correspondence:

University of Kirikkale
 Faculty of Dentistry
 Department of Pediatric Dentistry
 71200-Kirikkale
 Turkey
dr.volkanarikan@kku.edu.tr

Arikan V, Cumaogullari O, Ozgul BM, Oz FT. Investigation of SOSTDC1 gene in non-syndromic patients with supernumerary teeth. Med Oral Patol Oral Cir Bucal. 2018 Sep 1;23 (5):e531-9.
<http://www.medicinaoral.com/medoralfree01/v23i5/medoralv23i5p531.pdf>

Article Number: 22520 <http://www.medicinaoral.com/>
 © Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
 eMail: medicina@medicinaoral.com

Indexed in:

Science Citation Index Expanded
 Journal Citation Reports
 Index Medicus, MEDLINE, PubMed
 Scopus, Embase and Emcare
 Indice Médico Español

Received: 13/04/2018
 Accepted: 05/07/2018

Abstract

Background: The etiology of supernumerary teeth is still unclear however heredity is believed to be a major factor and this idea was supported by several case reports. Recently, a relationship between supernumerary tooth formation and deficiency of Uterine Sensitization Associated Gene-1 (*Usag-1*), a rat gene that is expressed in sensitized endometrium, was reported in mice. The human homolog gene for *Usag-1*, Sclerostin Domain Containing 1 (SOSTDC1), shows 85% identity with mouse *Usag-1*. The present study aimed to investigate *SOSTDC1* coding regions in non-syndromic patients with one or more supernumerary teeth.

Material and Methods: Twenty-five non-syndromic patients (21 male and 4 female) aged 5-15 years, with one or more supernumerary teeth were included in the study. Saliva samples were collected from patients and DNA samples were isolated and analyzed using PCR.

Results: Eight phenotypes of supernumerary tooth formation were observed in the study. From the DNA analysis, 2 novel and 3 previously identified sequence alterations were identified however, in investigating the *Usag-1* homolog *SOSTDC1* gene, the present study could not find any phenotype-genotype relationship.

Conclusions: There are many *SOSTDC1* homolog genes in the human genome and future studies should investigate these candidate genes. Also studies in larger case groups including family members may reveal the hereditary pattern.

Key words: Genetics, *Usag-1*, mesiodens, DNA sequencing, pediatric dentistry, PCR.

Introduction

Teeth produced in greater numbers than the normal dental formula are referred to as supernumerary teeth (1). The etiology of supernumerary teeth is still unclear, although there are some theories regarding the mechanism of their formation, including genetic and environmental factors (2). While this anomaly is most commonly observed in the

premaxilla, it can be found everywhere in the dental arch, and can occur in both primary and permanent dentition (3,4). Its prevalence ranges from 0.5% to 3.8% in permanent dentition and from 0.3% to 1.9% in primary dentition (5-8).

Heredity is believed to be a major factor behind supernumerary tooth formation. It has been suggested that

supernumerary teeth may be associated with autosomal recessive heredity, with lower penetrance in females (9). However, a few case reports have also proposed a low frequency autosomal dominant inheritance of this phenotype (10-12). Several articles support the idea that genetic component is needed for development of supernumerary teeth (13-15).

Recently, a relationship between supernumerary tooth formation and deficiency of Uterine Sensitization Associated Gene-1 (*Usag-1*), a rat gene that is expressed in sensitized endometrium (16), was reported in mice (17-19). Previous studies have shown that *Usag-1* and its human orthologous (ectodin) binds, neutralizes and acts as an antagonist of morphogenetic proteins (BMP). It has also been reported to inhibit Wnt signaling (18,20,21) while it is well established that both BMP and Wnt play a role in tooth morphogenesis (22,23), making it is very likely that *Usag-1* has a role in supernumerary teeth formation.

The human homolog gene for *Usag-1*, Sclerostin Domain Containing 1 (*SOSTDC1*), shows 85% identity with mouse *Usag-1*. The gene is localized on 7p21.1 with two transcripts of 3 and 5 exons, coding protein products of 206 and 230 amino acids respectively.

This study is the first investigation of *SOSTDC1* in humans with supernumerary teeth phenotype. We aimed to investigate *SOSTDC1* coding regions in non-syndromic patients with one or more supernumerary teeth.

Material and Methods

Twenty-five non-syndromic patients who were recruited through Ankara University's Department of Pediatric Dentistry between January 2011 and January 2013 with one or more supernumerary teeth were included in the study. There were 21 male and 4 female patients, aged 5-15 years. Additional patient information is given in the Table 1. Ethical approval was received from the Institutional Review Board (144/4), and informed consent was obtained from all participants and their parents.

- DNA isolation

Samples were collected from patients' saliva. Before isolating saliva DNA, the samples were dissolved in 10 ml isotonic solution in 15 ml falcon tubes. The tubes were centrifuged at 400g for 10 minute. Afterwards, the pellets were incubated at 56°C for 10 min, in 180 µl dH₂O and 20 µl proteinase K (20mg/ml). DNA isolations were performed using QIAamp DNA Blood Mini Kit (Qiagen Inc.) kit according to the manufacturer's instructions. DNA samples were spectrophotometrically analyzed and stored at -200C.

- Polymerase Chain Reaction (PCR)

To amplify the SOSTC1 gene (ENST00000396652), 3 pairs of primers were designed for coding exons (Table 2). Optimum PCR condition was obtained with 10 pmol primer mix, 0.5 unit SuperHot Taq polymerase enzyme (Bioron Inc.), and 1X Buffer Complete (Bioron Inc.) in

25 µl total volume. For each PCR reaction, a 20-50 ng/µl DNA template was used. Primer annealing temperature was optimized to 60°C. Thermal cycling conditions were 95°C 10 min for 1 cycle, 95°C 45 sec, 60°C 45 sec, 72°C 45 sec for 35 cycles, and 72°C 10 min. The PCR product was loaded in 2% agarose gel. Amplicon sizes of the PCRs were 531 bp (exon 3), 354 bp (exon 4) and 597 bp (exon 5).

- PCR Purification and DNA Sequencing

All PCR products were purified by using NucleoFast® 96 PCR kit (Macherey-Nagel GmbH). DNA sequencing was performed by cycle sequencing in 20 µl total volume. Sequencing reactions were set with both forward and reverse primers. Purification of the sequencing reaction was performed with ZR-96 DNA Sequencing Clean-up Kit (Zymo Research Corp.) according to the manufacturer's recommendations. Capillary electrophoresis was performed by ABI 3130 capillary electrophoresis instrument (Applied Biosystems Inc.). Electrophoregrams were analyzed by using SeqScape 2.5.0 software (Applied Biosystems Inc.).

- Bioinformatics analysis

DNA sequence results of the patients were aligned to Ensembl Grch37 Homo sapience SOSTDC1 coding regions nucleotide sequences to determine alterations. Subsequently, alterations were compared to "NCBI/NIH dbSNP (The Short Genetic Variations Database) short variations catalogs Homo sapience dataset"(24). The effect of missense mutation alterations in protein were investigated with SIFT analysis (25,26).

Results

A total of 25 patients (21 male, 4 female) were examined, of which 9 had single supernumerary teeth while the rest had 2 or more. Among the patients with single supernumerary teeth, 7 had mesiodentes (Table 3). Eight phenotypes were observed in our cohort (Table 4).

From the DNA analysis, we identified 2 novel and 3 previously identified sequence alterations (Table 5). The three previously discovered SNPs (Single Nucleotide Polymorphisms) are rs6945425, rs67149353, rs143801072. For the first two of these, the Minor Allele Frequencies (MAF) were 0.1212 and 0.2084 respectively in the dbSNP database. rs6945425 (c.-48A>G) and rs67149353 (IVS4+9G>C) are intronic nucleotide substitutions with no functional consequences on the protein. According to dbSNP, these SNPs have not been related with any syndrome. On the other hand, rs143801072 (c.476A>G, N159S), which shows a very low frequency in other Caucasian populations (0.010), is a missense mutation causing a change from asparagine to serine. We found rs143801072 in only one patient with phenotype 1 in heterozygous state, showing that the Asp>Ser amino acid alteration was tolerated by the protein. This patient was also found to carry rs6945425 variation in homozygous state.

The 2 novel mutations we identified had heterozygous genotype in phenotypes 3 and 5. One of these alterations,

Table 1. DNA analysis and additional patient information.

Variation type:	known intronic variation	known intronic variation	novel deleterous variation	known missence variation	novel silent variation	
Patient number	rs6945425 (c.-48A>G)	rs671149353 (Ivs4+9G>C)	c.221-223delCGA (p.Leu73del)	rs143801072 (c.476A>G, p.Asn159Ser)	c.298C>T (p.Asn99Asn)	Phenotype Name
1	HET	HET	WT	WT	WT	Phenotype 5
2	MUT	HET	WT	WT	WT	Phenotype 5
3	HET	WT	WT	WT	WT	Phenotype 3
4	MUT	aWT	WT	WT	WT	Phenotype 5
5	MUT	HET	WT	WT	WT	Phenotype 8
6	MUT	WT	WT	WT	WT	Phenotype 3
7	MUT	MUT	WT	WT	WT	Phenotype 5
8	MUT	WT	WT	WT	WT	Phenotype 5
9	MUT	WT	WT	WT	WT	Phenotype 6
10	HET	HET	WT	WT	WT	Phenotype 8
11	MUT	WT	WT	WT	WT	Phenotype 8
12	MUT	WT	WT	HET	WT	Phenotype 1
13	MUT	WT	WT	WT	WT	Phenotype 3
14	MUT	HET	WT	WT	WT	Phenotype 3
15	MUT	WT	WT	WT	HET	Phenotype 3
16	HET	WT	WT	WT	WT	Phenotype 7
17	MUT	HET	WT	WT	WT	Phenotype 5
18	MUT	WT	WT	WT	WT	Phenotype 5
19	HET	WT	WT	WT	WT	Phenotype 2
20	MUT	HET	WT	WT	WT	Phenotype 5
21	MUT	HET	WT	WT	WT	Phenotype 4
22	MUT	HET	WT	WT	WT	Phenotype 5
23	MUT	HET	WT	WT	WT	Phenotype 3
24	MUT	HET	HET	WT	WT	Phenotype 5
25	HET	HET	WT	WT	WT	Phenotype 3

*HET: Heterozygous genotype, HOM: Homozygous genotype, WT: Wild type.

Table 2. Sequences of primers used of the amplification of SOSTC1 gene.

Gene	Exon	Orientation	Sequence (5'->3')
USAG1	3	Forward	GGCAATTTGTATACCAAGCTCCTCC
USAG1	3	Reverse	ATTCTACAGGAATGTGAGCTAATGCTACCAG
USAG1	4	Forward	GTTTTAACTTTCACGAAGCTGGTTGC
USAG1	4	Reverse	GCTTAAGGGGAACGTGATAGTTGTGG
USAG1	5	Forward	CTATCATTTCCTATTATTTTGTTCATTGC
USAG1	5	Reverse	GCAGTGGCAGGCTTGAGTCTTCC

Table 3. Distribution on number, region and types of supernumerary teeth.

Patient	Gender	Number of extra teeth	Region and Type
1	M	2	Bilateral maxillary centrals
2	M	2	Bilateral maxillary centrals
3	M	1	Mesiodens
4	M	2	Bilateral maxillary centrals
5	M	2	Unilateral maxillary primary and permanent centrals
6	M	1	Mesiodens
7	M	2	Bilateral maxillary centrals
8	M	2	Bilateral maxillary centrals
9	M	5	Bilateral maxillary central and laterals
10	M	2	Unilateral maxillary primary and permanent centrals
11	M	2	Unilateral maxillary primary and permanent centrals
12	F	6	Bilateral maxillary centrals and bilateral mandibular 1st and 2nd premolars
13	M	1	Mesiodens
14	M	1	Mesiodens
15	M	1	Mesiodens
16	M	1	Unilateral maxillary lateral
17	M	2	Bilateral maxillary centrals
18	F	2	Bilateral maxillary centrals
19	M	1	Unilateral mandibular premolar
20	F	2	Bilateral maxillary centrals
21	M	5	Bilateral maxillary centrals and mandibular premolars
22	M	2	Bilateral maxillary centrals
23	F	1	Mesiodens
24	M	2	Bilateral maxillary centrals
25	M	1	Mesiodens

Table 4. Phenotypic variations.

Name	Phenotype	n
Phenotype 1	Bilateral maxillary centrals and bilateral mandibular 1st and 2nd premolars	1
Phenotype2	Unilateral mandibular premolar	1
Phenotype3	Mesiodens	7
Phenotype4	Bilateral maxillary centrals and mandibular premolars	1
Phenotype5	Bilateral maxillary centrals	10
Phenotype6	Bilateral maxillary central and laterals	1
Phenotype7	Unilateral maxillary lateral	1
Phenotype8	Unilateral maxillary primary and permanent centrals	3

Table 5. Identified alterations in our study.

Nucleotide change and genotypes	Rs no	Number	% Presence	Localization on the gene	Effect on protein
c.-48A>G_	rs6945425				
AA		0	0	Intron 2	-
AG		20	76,9	Intron 2	-
GG		6	23,1	Intron 2	-
IVS4+9G>C_	rs67149353		46,2		-
GG		12		Intron 4	
GC		13	50	Intron 4	-
CC		1	3,8	Intron 4	-
c.221-223delCGA					
WT		25	96,2	Exon 4	-
WT/3del		1	3,8	Exon 4	p.Thr74del
3del/3del		0	0	Exon 4	=
c.476A>G_	rs143801072				
AA		25	96,2	Exon 5	-
AG		1	3,8	Exon 5	p.Asn159Ser
GG		0	0	Exon 5	=
c.298C>T					
CC		25	96,2	Exon 5	-
CT		1	3,8	Exon 5	p.Asn99Asn
TT		0	0	Exon 5	=

c.221-223delCGA, was an in-frame deletion causing one tyrosine amino acid to be deleted in SOSTDC1 (Table 1). This was detected in 1 in 10 patients with phenotype 5. The other novel nucleotide alteration, c.298C>T, is a synonymous mutation that does not change asparagine amino acid at residue 99. This substitution was identified in 1 in 7 cases with phenotype 3.

Discussion

The etiology behind the formation of supernumerary teeth still remains unknown. However, several theories have been investigated previously. One of these theories was dichotomy, which claims that the developing tooth bud may be divided to form a supernumerary tooth (27). Hyperactivity of the dental lamina has also been suggested as a possible factor behind the formation of supernumerary teeth (4).

Various genes (*RUNX2*, *PLOD*, *EVC*, *GLA*, *APC*, *NEMO*) have been associated with supernumerary teeth formation in several syndromes, such as Cleidocranial dysplasia, Ehlers-Danlos Type IV, Ellis-Van Creveld, Fabry disease, Familial adenomatous polyposis, and Incontinentia pigmenti (13,28,29). As previously noted, our study focused on non-syndromic supernumerary tooth formation.

In the last two decades, several studies conducted on mutation induced mice in *Usag-1*, *Gas1*, *Eda*, *Spry 2*, *Spry 4*, and *Pax 6* resulted in supernumerary tooth formation (14, 15,17,30-33).

Even though the etiology behind supernumerary teeth cannot be clearly determined, studies have shown that cell cycle related pathways like WNT, MAPK/ERK, and PI3K/AKT/Mtor are involved in supernumerary tooth formation. It is well known that BMP and Wnt are key molecules controlling tooth morphogenesis (22). BMP is known to regulate embryonic development in all animals, being present in practically all tissues and organs. *Usag-1* (also known as *Ectodin*, *Sostdc1* or *Wise*), which is expressed in the epithelium and mesenchyme of the developing tooth germ, encodes a secreted BMP-inhibitor (21, 34). Recently, Murashima-Suginami *et al.* have shown that *Usag-1* abrogation in mice resulted in the survival of rudiment incisors and formation of supernumerary teeth (18). In a different study, their team also found that BMP signaling increases in *Usag-1* deficient mice since this gene is an antagonist of BMPs, which results in the formation of supernumerary teeth (19). According to Kiso *et al.*, as a result of *Usag-1* gene deficiency, due to lack of apoptotic elimination, odontogenic mesenchymal cells were retained in mice, while the interaction between *Bmp-7* and *Usag-1* had a role in the formation of supernumerary organs (35). Kassai *et al.* also reported supernumerary tooth formation in *Usag-1* (which they named *Ectodin*) deficient mice (17). In this study, we investigated *Usag-1* gene homolog *SOSTDC1* in 25 patients with at least one supernumerary tooth without any syndromes. The majority (84%) of our study

group were male patients. We detected 2 novel and 3 previously identified nucleotide alterations (Table 4).

The known variations rs6945425 (c.-48A>G) and rs67149353 (IVS4+9G>C) are intronic nucleotide substitutions with no functional consequences on the protein whereas, as previously noted, rs143801072 (c.476A>G) is a missense mutation (p.Asn159Ser). This residue (p.Asn159Ser) is found to be conserved in many species (Table 6). MAF is 0.01 among Caucasian populations. This residue lies within the evolutionarily conserved C-terminal cystine knot-like (CTCK) domain. As in other hereditary diseases, such as Bardet Biedl and Meckel Syndrome, hereditary non-syndromic supernumerary phenotype seems to be multigenic. Thus, this rare SNP in a functional domain in a very highly conserved residue may cause supernumerary teeth formation in just 1 in 10 patients with phenotype 1. It is important to emphasize that phenotype 1 was found in only one patient with rs143801072 heterozygous variation. However, this patient, for whom we lacked the family history, was the only case where we identified a potentially deleterious heterozygous mutation. It is therefore not possible to fully establish the relationship between supernumerary tooth formation, phenotype 1 and rs143801072. Besides, rs6945425 homozygous genotype was also detected in this patient, along with other 20 patients. Further investigations are therefore needed to elucidate the functional consequences of this missense mutation.

We also identified 2 novel mutations, c.221-223delCGA and c.298C>T. One patient was heterozygous for the c.221-223delCGA alteration, which causes an in-frame deletion of tyrosine at residue 74 in the *SOSTDC1* protein (Table 1). Further functional assays need to be conducted in order to establish whether this deletion impairs protein function. We evaluated probable functional consequences of this amino acid deletion in silico using three dimensional structure analysis tools (Coils regions, Domain linker prediction, Helical context). The analyses showed that the deletion of tyrosine at residue 74 did not have any structural or functional effect on the protein. Besides, this residue is not conserved across species. According to the sift analysis, this variation can be tolerated without affecting protein function.

The other novel nucleotide variation, c.298C>T, is a silent mutation that does not change asparagine at residue 99. Thus, this substitution has no functional impact on *SOSTDC1*.

In investigating the *Usag-1* homolog *SOSTDC1* gene, we could not find any phenotype-genotype relationship. According to our in silico analysis, there are many *SOSTDC1* homolog genes in the human genome (Table 7). Future studies should investigate these candidate genes within the same study group. It would also be useful to enlarge the case group, and include family members, which may reveal the hereditary pattern.

Table 6. Comparison of SOSTDC1 protein in different species with protein domains and variation points.

Homo sapiens_Q6X4U4-2	1	ML---PPAIHFYLLPLACILMKSCAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFSTGLDRNRTESEITFQGN	77
Pan troglodytes_H2RCY3	1	ML---PPAIHFYLLPLACILMKSCAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFSTGLDRNRTESEITFQGN	77
Gorilla gorilla_G3R2X4	1	ML---PPAIHFYLLPLACILMKSCAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFSTGLDRNRTESEITFQGN	77
Mus musculus_Q9CQN4	1	ML---PPAIHLSLIPLLCILMRNCLAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFSTGLDRN-----	68
Rattus norvegicus_Q642G2I	1	ML---PPAIHLSLIPLLCILMKNCLAFKNDATEILYSHVVKVPSAHPSSNSTLNOARNGGRHFSTGLDRN-----	68
Canis lupus familiaris_F1PUT3	1	ML---PPAIHFYLLPLACILMKSCAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFSTGLDRNSNKILGSSQN	77
Gallus gallus_Q6VYA3	1	ML---LSAIHFYGLLACTFTRSYAFKNDATEILYSHVVKVPVPAHPSSNSTLNOARNGGRHFYAGTGSDRN-----	68
Danio rerio_F1QVR9	1	MYINAPESCNFMV--LFCFLIRSQLTKINDATEIFYSHVSPVQ-DAQSNASLNRAASGRGFSTH--DRE-----	68
Homo sapiens_Q6X4U4-2	78	YFWLFPFGAFRLQLQEARVQVGC CRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	157
Pan troglodytes_H2RCY3	78	YFWLFPFGAFRLQLQEARVQVGC CRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	157
Gorilla gorilla_G3R2X4	78	YFWLFPFGAFRLQLQEARVQVGC CRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	157
Mus musculus_Q9CQN4	69	-----SRVQVGC CRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	133
Rattus norvegicus_Q642G2I	69	-----SRVQVGC CRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	133
Canis lupus familiaris_F1PUT3	78	HLWVFPFGAFLGQMDEARVQV SCRELRSTKYISDGGQCTSISPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	157
Gallus gallus_Q6VYA3	69	-----NRVQVGC CRELRSTKYISDGGQCTSINPLKELVCAGECLPLPLPNWIGGGYGTKYWSRRSSQEWRC	133
Danio rerio_F1QVR9	67	-----RIPVGC CRELRSTKYISDGGQCTSINPVKELVCTGQCLPAQMLPNWIGG-YGKKSWNRRNSQEWRC	129
Homo sapiens_Q6X4U4-2	158	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SMS---PAKPVQHHRRERKRASKSKSHSMS	230
Pan troglodytes_H2RCY3	158	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SMS---PAKPVQHHRRERKRASKSKSHSMS	230
Gorilla gorilla_G3R2X4	158	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SMS---PAKPVQHHRRERKRASKSKSHSMS	230
Mus musculus_Q9CQN4	134	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SVS---PAKPAQHHRERKRASKSKSHLS	206
Rattus norvegicus_Q642G2I	134	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SVS---PAKPAQHHRERKRASKSKSHLS	206
Canis lupus familiaris_F1PUT3	158	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE SMS---PAKPAQHHRERKRASKSKSHLS	230
Gallus gallus_Q6VYA3	134	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYTRQHNESSHNFE GTS---QAKPVQHHKERKRASKSKSHSTS	206
Danio rerio_F1QVR9	130	VNDKTRTRQRIQLQCDDGSTRTYKITVVTACKCKRYSRQHNESSGVKSEGYSHSQIKTEKQSGHODRKLKLSLELTLI	206

p.Thr74del

CTCK domain

p.Asn159Se

Table 7. SOSTDC1 90% and more homologue genes in the human genome (Blast-Ensembl tool).

Chromosome	Overlapping Gene(s)	Query start	Query end	Length (Sequence)	Score	E-val	%ID (Alignment)
7	SOSTDC1	28607	69100	40494	78394	0.00E+00	100
7	GS1-166A23.1	28607	69100	40494	78394	0.00E+00	100
4	HERC3	44028	44322	295	516	1.30E-146	94.24
1	AGBL4	44026	44320	295	510	1.10E-144	92.88
2	LPIN1	44196	44508	314	521	3.70E-148	92.68
5	RP11-541P9.3	43956	44322	367	616	1.10E-176	92.37
X	MAP3K15	44036	44443	408	683	5.90E-197	92.16
3	RP11-23D24.2	44031	44508	479	792	9.90E-230	92.07
3	ROBO1	43960	44446	487	815	1.80E-236	91.99
7	NOD1	44026	44428	403	668	1.80E-192	91.81
3	STXBP5L	44084	44508	426	703	6.20E-203	91.78
6	LAMA4	43976	44508	534	883	5.10E-257	91.76
13	RP11-141M1.3	44024	44446	423	698	2.20E-201	91.73
3	TBC1D5	43949	44299	351	588	2.50E-168	91.45
1	DNAH14	44026	44350	325	534	6.70E-152	91.38
1	DNAH14	44053	44446	394	652	1.70E-187	91.37
8	RGS22	44022	44446	425	705	1.90E-203	91.29
6	SAMD3	43956	44466	512	837	3.80E-243	91.21
7	PPP1R9A	43976	44503	529	851	2.30E-247	90.93
8	RP11-770E5.1	43977	44428	452	740	5.00E-214	90.93
4	CENPE	44026	44508	484	779	9.70E-226	90.91
3	WDR49	43959	44363	408	655	1.50E-188	90.69
10	FRMD4A	43956	44508	554	890	4.40E-259	90.43
6	ADGRB3	44022	44508	489	786	6.30E-228	90.39
1	KCNK2	44032	44391	360	577	7.30E-165	90.28
4	RP11-8L2.1	43977	44499	524	836	6.10E-243	90.27
4	RP11-103J17.1	44036	44446	411	664	5.20E-191	90.02
3	LINC00578	44025	44503	480	759	7.40E-220	90

References

1. Ferrer-Padro E, Prats-Armengol J, Ferrer-Amat E. A descriptive study of 113 unerupted supernumerary teeth in 79 pediatric patients in Barcelona. *Med Oral Patol Oral Cir Bucal.* 2009;14:E146-52.
2. Rajab LD, Hamdan MA. Supernumerary teeth: review of the literature and a survey of 152 cases. *Int J Paediatr Dent.* 2002;12:244-54.
3. Celikoglu M, Kamak H, Oktay H. Prevalence and characteristics of supernumerary teeth in a non-syndrome Turkish population: associated pathologies and proposed treatment. *Med Oral Patol Oral Cir Bucal.* 2010;15:e575-8.
4. Anthonappa RP, Omer RS, King NM. Characteristics of 283 supernumerary teeth in southern Chinese children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008;105:e48-54.
5. Arıkan V, Özgül BM, Fırdevs TO. Prevalence and characteristics of supernumerary teeth in a child population from Central Anatolia - Turkey. *Oral Health Dent Manag.* 2013;12:269-72.
6. Backman B, Wahlin YB. Variations in number and morphology of permanent teeth in 7-year-old Swedish children. *Int J Paediatr Dent.* 2001;11:11-7.
7. Fardi A, Kondylidou-Sidira A, Bachour Z, Parisis N, Tsirlis A. Incidence of impacted and supernumerary teeth-a radiographic study in a North Greek population. *Med Oral Patol Oral Cir Bucal.* 2011;16:e56-61.
8. Zhu JF, Marcushamer M, King DL, Henry RJ. Supernumerary and congenitally absent teeth: a literature review. *J Clin Pediatr Dent.* 1996;20:87-95.
9. Niswander JD, Sujaku C. Congenital Anomalies of Teeth in Japanese Children. *Am J Phys Anthropol.* 1963;21:569-74.
10. Batra P, Duggal R, Parkash H. Non-syndromic multiple supernumerary teeth transmitted as an autosomal dominant trait. *J Oral Pathol Med.* 2005;34:621-5.
11. Orhan AI, Ozer L, Orhan K. Familial occurrence of nonsyndromal multiple supernumerary teeth. A rare condition. *Angle Orthod.* 2006;76:891-7.
12. Wang XX, Zhang J, Wei FC. Autosomal dominant inheritance of multiple supernumerary teeth. *Int J Oral Maxillofac Surg.* 2007;36:756-8.
13. Fleming PS, Xavier GM, DiBiase AT, Cobourne MT. Revisiting

- the supernumerary: the epidemiological and molecular basis of extra teeth. *Br Dent J.* 2010;208:25-30.
14. Kangas AT, Evans AR, Thesleff I, Jernvall J. Nonindependence of mammalian dental characters. *Nature.* 2004;432:211-4.
15. Anthonappa RP, King NM, Rabie AB. Aetiology of supernumerary teeth: a literature review. *Eur Arch Paediatr Dent.* 2013;14:279-88.
16. Simmons DG, Kennedy TG. Uterine sensitization-associated gene-1: a novel gene induced within the rat endometrium at the time of uterine receptivity/sensitization for the decidual cell reaction. *Biol Reprod.* 2002;67:1638-45.
17. Kassai Y, Munne P, Hotta Y, Penttila E, Kavanagh K, Ohbayashi N, et al. Regulation of mammalian tooth cusp patterning by ectodin. *Science.* 2005;309:2067-70.
18. Murashima-Suginami A, Takahashi K, Kawabata T, Sakata T, Tsukamoto H, Sugai M, et al. Rudiment incisors survive and erupt as supernumerary teeth as a result of USAG-1 abrogation. *Biochem Biophys Res Commun.* 2007;359:549-55.
19. Murashima-Suginami A, Takahashi K, Sakata T, Tsukamoto H, Sugai M, Yanagita M, et al. Enhanced BMP signaling results in supernumerary tooth formation in USAG-1 deficient mouse. *Biochem Biophys Res Commun.* 2008;369:1012-6.
20. Itasaki N, Jones CM, Mercurio S, Rowe A, Domingos PM, Smith JC, et al. Wise, a context-dependent activator and inhibitor of Wnt signalling. *Development.* 2003;130:4295-305.
21. Laurikkala J, Kassai Y, Pakkasjarvi L, Thesleff I, Itoh N. Identification of a secreted BMP antagonist, ectodin, integrating BMP, FGF, and SHH signals from the tooth enamel knot. *Dev Biol.* 2003;264:91-105.
22. Peters H, Balling R. Teeth. Where and how to make them. *Trends Genet.* 1999;15:59-65.
23. Stockton DW, Das P, Goldenberg M, D'Souza RN, Patel PI. Mutation of PAX9 is associated with oligodontia. *Nat Genet.* 2000;24:18-9.
24. Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, Smigielski EM, et al. dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* 2001;29:308-11.
25. Ng PC, Henikoff S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res.* 2003;31:3812-4.
26. Ng PC, Henikoff S. Predicting the effects of amino acid substitutions on protein function. *Annu Rev Genomics Hum Genet.* 2006;7:61-80.
27. Khambete N, Kumar R. Genetics and presence of non-syndromic supernumerary teeth: A mystery case report and review of literature. *Contemp Clin Dent.* 2012;3:499-502.
28. Lubinsky M, Kantaputra PN. Syndromes with supernumerary teeth. *Am J Med Genet A.* 2016;170:2611-6.
29. Subasioglu A, Savas S, Kucukyilmaz E, Kesim S, Yagci A, Dunder M. Genetic background of supernumerary teeth. *Eur J Dent.* 2015;9:153-8.
30. Ohazama A, Haycraft CJ, Seppala M, Blackburn J, Ghafoor S, Cobourne M, et al. Primary cilia regulate Shh activity in the control of molar tooth number. *Development.* 2009;136:897-903.
31. Kaufman MH, Chang HH, Shaw JP. Craniofacial abnormalities in homozygous Small eye (*Sey/Sey*) embryos and newborn mice. *J Anat.* 1995;186 (Pt 3):607-17.
32. Peterkova R, Churava S, Lesot H, Rothova M, Prochazka J, Peterka M, et al. Revitalization of a diastemal tooth primordium in *Spry2* null mice results from increased proliferation and decreased apoptosis. *J Exp Zool B Mol Dev Evol.* 2009;312B:292-308.
33. Klein OD, Minowada G, Peterkova R, Kangas A, Yu BD, Lesot H, et al. Sprouty genes control diastema tooth development via bidirectional antagonism of epithelial-mesenchymal FGF signaling. *Dev Cell.* 2006;11:181-90.
34. Yanagita M, Oka M, Watabe T, Iguchi H, Niida A, Takahashi S, et al. USAG-1: a bone morphogenetic protein antagonist abundantly expressed in the kidney. *Biochem Biophys Res Commun.* 2004;316:490-500.
35. Kiso H, Takahashi K, Saito K, Togo Y, Tsukamoto H, Huang B, et al. Interactions between BMP-7 and USAG-1 (uterine sensitization-associated gene-1) regulate supernumerary organ formations. *PLoS One.* 2014;9:e96938.

Acknowledgements

This study was funded by Ankara University Faculty of Dentistry Scientific Research Projects Coordination Unit.

The authors would like to thank Prof.Dr.Hilal Özdağ from Ankara University Biotechnology Institute for her generous aid and guidance throughout the study.

Conflict of interest

All authors of the manuscript declare that they have no conflicts of interest.