



## Stress Responses of Stallions During Transport Period: The Variations of Complete Blood Count and Serum Biochemistry\*

Ali Evren HAYDARDEDEOĞLU<sup>1</sup>, Buğrahan Bekir YAĞCI<sup>2</sup>, Kerem URAL<sup>3</sup>, Hasan GÜZELBEKTEŞ<sup>4</sup>

1. Aksaray University, Faculty of Veterinary Medicine, Department of Internal Medicine, Aksaray, TURKEY.
2. Kırıkkale University, Faculty of Veterinary Medicine, Department of Internal Medicine, Kırıkkale, TURKEY.
3. Adnan Menderes University, Faculty of Veterinary Medicine, Department of Internal Medicine, Aydın, TURKEY.
4. Selçuk University, Faculty of Veterinary Medicine, Department of Internal Medicine, Konya, TURKEY.

Geliş Tarihi/Received	Kabul Tarihi/Accepted	Yayın Tarihi/Published
19.08.2016	21.02.2017	30.10.2017

**Abstract:** In the present study the objective was to determine the effect of one transport journey on some stress indicators in horses. 11 adult stallion (all stallion) age 3-12 years including 7 Thoroughbred, 2 Arabian, 1 Dutch Warmblood, and 1 Friesian were enrolled. White blood cell counts and other relevant parameters were determined by an automated cell counter before and after transportation. Serum activities of relevant biomarkers were determined by use of an autoanalyzer. Serum cortisol level was detected by a microplate immunoassay technique. Sodium (Na), Potassium (K) and Chlor (Cl) and Calcium (Ca) levels were also measured by use of autoanalyser. Mean Platelet (PLT), and Mean corpuscular hemoglobin concentration (MCHC) White Blood Cell (WBC) counts did not exhibit a variation for healthy horses during the study whereas RBC, Hb and Hct levels exhibited significant increases after transportation ( $P < 0.05$ ). Mean daily glucose concentration showed an increase after transportation ( $P < 0.05$ ). Mean total protein, albumin, AST and CK levels presented increases on day 2, after transportation ( $P < 0.05$ ). Mean blood urea nitrogen, creatinine, total bilirubin, ALT, ALP, LDH and GGT did not show significant changes throughout the study. Mean Sodium (Na), Potassium (K) and Chlor (Cl) levels showed alterations after transportation ( $P < 0.05$ ), whereas Calcium (Ca) levels showed significant decreases after transportation ( $P < 0.05$ ).

**Keywords:** Biochemistry, Hematology, Stallion, Stress, Transport.

## Uzun Transporta Bağlı Aygırların Strese Tepkileri: Tam Kan Sayımı ve Serum Biyokimya Varyasyonları

**Öz:** Bu çalışmada, transportun atlarda bazı stres göstergelerine etkisini belirlemek amaçlanmıştır. 7 Safkan, 2 Arap, 1 Hollanda Yarım Kan ve 1 Fresyan olmak üzere 3-12 yaş arası 11 olgun at (aygır) kayıt edildi. Transport öncesi ve sonrasında, otomatik kan sayımı ile beyaz kan hücresi sayımı ve diğer ilgili parametreler belirlendi. İlgili biyolojik belirteçlerin serum aktiviteleri bir otoanalizör kullanılarak belirlendi. Na, K, Cl ve Ca düzeyleri de otoanalizör kullanılarak ölçüldü. Ortalama Platelet (PLT) ve Ortalama Hemoglobin Konsantrasyonu (MCHC) Total Lökosit (WBC) sayıları çalışma süresince sağlıklı atlar için bir farklılık göstermezken, Alyuvarlar (RBC), Hemoglobin (Hb) ve Hematokrit (Hct) düzeyleri ulaşımdan sonra anlamlı artışlar sergiledi ( $P < 0.05$ ). Günlük ortalama glikoz konsantrasyonu transporttan sonra artış gösterdi ( $P < 0.05$ ). Ortalama total protein, albümin, Aspartat aminotransferaz (AST) ve Kreatinin kinaz (CK) düzeyleri, transportun ardından 2. günde artmaktadır ( $P < 0.05$ ). Ortalama kan üresi azotu, kreatinin, total bilirubin, Alanin aminotransferaz (ALT), Alkalen fosfataz (ALP), Laktat dehidrogenaz (LDH) ve Gama-gulutamil Transferaz (GGT), çalışma boyunca önemli değişiklikler göstermedi. Transporttan sonra ortalama Sodyum (Na), Potasyum (K) ve Klor (Cl) düzeyleri değişti ( $P < 0.05$ ), nakil sonrası Kalsiyum (Ca) düzeyleri anlamlı düşüş gösterdi ( $P < 0.05$ ).

**Anahtar Kelimeler:** Aygır, Biyokimya, Hematoloji, Stres, Transport.

<sup>1</sup>Ali Evren HAYDARDEDEOĞLU

Aksaray University, Faculty of Veterinary Medicine, Department of Internal Medicine, Aksaray, TURKEY.  
e-mail: ahaydardedeoglu@aksaray.edu.tr

\*This article was presented as poster in the 32nd WVA congress, 2015-Istanbul, Turkey.

## INTRODUCTION

Horses are transported for several reasons comprising sport (showing and racing) (1,2,3), breeding and slaughter for several times throughout their lives (4). It has been well recognized that in horses transport is a general stressor leading to salmonellosis (5), respiratory disease (6,7,8), elevated heart rates (9,10), and changes in plasma ascorbic acid (11), serum cortisol concentration (9,10,11,12,13) and muscle enzyme activities (13,14,15). Elevated sodium and blood glucose concentrations have also been detected in slaughtering horses (12,13,14,15,16).

Up to date the transportation of horses by road received attention, but limited scientific research has been (4) focused on both hematological and serum biochemical changes. Therefore the aim of this study was to determine the effect of one transport journey on some stress indicators in horses.

## MATERIALS and METHODS

### Horses and Study Design

Eleven adult stallion aged 3-12 years including 7 Thoroughbred, 2 Arabian, 1 Dutch Warmblood, and 1 Friesian were enrolled. All horses have had prior experience of transport; indeed 2 weeks prior to the study was performed, all horses were loaded and unloaded from the van in an attempt to ensure that all horses loaded without hesitation. This study was conducted in accordance with the ethical guidelines of animal experiments.

A study design consisted of one 19 h transport period from Nevsehir to Antalya, with a total of 867 km travelled. On day 0, prior to study horses were housed individually in their natural boxes and physiological data were collected to obtain baseline values. On day 1 horses were loaded to a commercial van. Travel commenced for 19 hour period. Horses were unloaded on day 2 and housed individually in boxes for post-transport period. Water was always available to the horses in their boxes.

A commercial van was designed to carry out a total of 11 stallions and was pulled by a tractor.

Horses were randomly enrolled in groups parallel to the length of the van, as described previously (21). Five horses faced the direction of the journey, and six other faced the opposite direction. Bedding (straw) was placed on the floor for secure footing. Windows were allowed to open fully for fresh air. Travel was initiated at early morning and continued for 19 h.

### Laboratory Analyses

White blood cell counts and were determined by an automated cell counter (Abacus Junior Vet 5, Hungary). Serum activities of Alanine aminotransferase (ALT), Aspartate aminotransferase (AST), Alkaline phosphatase (ALP), Creatinine kinase (CK), Lactate dehydrogenase (LDH), Gama-glutamyl transferase (GGT) and glucose, Blood Urea Nitrogen (BUN), creatinine, total protein, albumine, total bilirubine levels were determined by use of an autoanalyzer (Spotchem ez sp-4430, Arkray, Japan). Serum cortisol levels and electrolyte analysis [Na, K, Cl and Ca], were detected by a microplate immunoassay technique using autoanalyzer in a private commefical lab.

### Statistical Analyses

The effects of transportation and measurements taken after transport were evaluated using means for day 1 and 2, respectively, using Paired Samples t-test. Significance was declared whenever  $P < 0.05$ .

## RESULTS

### Blood Measurements

The WBC counts did not exhibit a variation and remained within the reference range ( $5.5$  to  $14.3 \times 10^3/_L$ ) for healthy horses during the study. Mean PLT and MCHC did not show significant levels (Table 1). RBC, Hb and Hct levels exhibited variations with increases after transportation ( $P < 0.05$ ). MCHC exhibited variations with slight decrease after transportation ( $P < 0.05$ ) (Table 1).

**Table 1.** Mean haematological values ( $\pm$  SD) of measurements and P-values of comparisons between day 0 (d 0) and following a 19-h posttransport period (d 2).

**Tablo 1.** 0. Hematolojik ölçüm değerlerinin ortalaması ( $\pm$  SD) ve 0.(d 0) gün ile 19 saatlik bir nakliye sonrası dönem (d 2) arasındaki karşılaştırmaların P-değerleri.

	N	O. hour	19. hour	SD	P	
WBC ( $\times 10^3/\mu\text{L}$ )	11	7.67	7.73	0.815	0.809	
RBC ( $\times 10^6/\mu\text{L}$ )	11	7.77	12.82	2.005	0.000	***
HB (g/dL)	11	11.91	16.93	1.415	0.000	***
HCT (%)	11	40.54	50.79	1.768	0.000	***
MCV (fL)	11	46.40	40.88	5.605	0.006	***
MCHC (g/dL)	11	33.39	33.50	1.877	0.845	
PLT ( $\times 10^6/\mu\text{L}$ )	11	150.25	147.92	6.945	0.269	

### Serum Metabolites and Enzymes

Mean daily glucose concentration showed an increase after transportation ( $P < 0.05$ ). (Table 2). Mean total protein, albumine, AST and CK levels showed increases on day 2, after transportation ( $P < 0.05$ ). Neither serum concentrations of blood urea nitrogen nor of creatinin exhibited variation or difference ( $P > 0.05$ ) between days 1 and 2. Besides mean values of total bilirubine, ALT, ALP, LDH and

GGT did not show significant changes throughout the study (Table 2).

Mean Na, K and Cl levels showed variations with increases after transportation ( $P < 0.05$ ), whereas Ca levels showed significant decreases after transportation ( $P < 0.05$ ). Mean serum cortisol levels showed significant increases after transportation ( $P < 0.05$ ) (Table 2).

**Table 2.** Mean serum biochemical and metabolite values ( $\pm$  SD) of measurements and P-values of comparisons between day 0 (d 0) and following a 19-h posttransport period (d 2). Daily means (SD) and P values.

**Tablo 2.** Ortalama serum biyokimyasal ve metabolit değerleri ( $\pm$  SD) ve 0. gün (d 0) ile 19 saat sonrası nakliye sonrası dönem (d 2) arasındaki karşılaştırmaların P değerleri. Günlük ortalama (SD) ve P değerleri.

	N	O. hour	19. hour	SD	P	
Glucose (mg/dL)	11	89.83	111.83	5.846	0.000	***
Blood Urea Nitrogen (mg/dL)	11	14.33	14.50	1.800	0.754	
Creatinin (mg/dL)	11	1.11	1.14	0.143	0.438	
Total Protein (g/dL)	11	6.25	7.53	0.407	0.000	***
Albumine (g/dL)	11	3.20	3.82	0.332	0.000	***
Total Bilirubine (mg/dL)	11	1.34	1.33	0.144	0.845	
ALT (IU/L)	11	13.75	14	2.800	0.763	
AST (IU/L)	11	311.17	396.67	40.218	0.000	***
ALP (IU/L)	11	188.08	187	6.815	0.593	
CK (IU/L)	11	12.33	28.84	3.091	0.000	***
LDH (IU/L)	11	271.17	270.25	9.209	0.737	
GGT (IU/L)	11	13	13.33	1.614	0.489	
Na <sup>+</sup> (mmol/L)	11	138.17	149.50	1.442	0.000	***
K <sup>+</sup> (mmol/L)	11	3.07	4.49	0.362	0.000	***
Cl <sup>-</sup> (mmol/L)	11	102.75	119.17	4.420	0.000	***
Ca <sup>++</sup> (mmol/L)	11	3.09	2.69	0.175	0.000	***
Cortisole ( $\mu\text{g/dL}$ )	11	1.65	2.90	0.215	0.000	***

## DISCUSSION and CONCLUSION

One of the most important problem is that the response to road transport equine practice worldwide (24). As a consequence several immunological, physiological, and metabolic alterations might prone horses to elevated susceptibility to disorders (diarrhea, colics, pneumonia laminitis and trauma) (24,25). Transportation is composed of several psychological stimuli, resulting with stress disrupting metabolism and homeostasis of affected animals (26,27,28).

Transportation includes various stressors comprising loading, unloading and penning in a novel and unfamiliar environment and confinement along with motion, vibrations, changes in climatic conditions, inadequate ventilation and food and water deprivation. Animals react and addict respond to challenges in their environment by various interacting mechanisms including behavioural, haematological, physiological and neuro-hormonal parameters. Age, sex and physiological conditions may also have influence on the behaviour of animals during handling and transport (17,18,19).

In a prior study analyzing relationships between transportation of 24 cold-blooded female horses forwarded to both long and short distances and an inflammatory condition [acute phase proteins, oxidative stress and muscle injury. Plasma and serum were withdrawn prior to transportation (T0), immediately after transportation (T1) and finally during slaughter (T2). Fibrinogen was elevated in all studied groups especially in fillies after long distance transportation, ( $205 \pm 7.07$  mg/dl at T0;  $625 \pm 35.35$  mg/dl at T1, and  $790 \pm 14.14$  mg/dl at T2). Malondialdehyde levels rose at T1 and reached the maximal level during T2. CK activity was also increased at T1 in younger horses, According to the latter study an intensified acute phase response along with oxidative stress and muscle trauma indicated an inflammatory condition (24). In the present study although the present authors did not evaluate direct biomarkers of stress or oxidative status, cortisol was markedly increased in horses

following transportation indicating relative stress as was also determined (25,29).

Significantly higher levels of GOT, GPT, LDH and ALT in contrast to the basal values were reported for sedentary horses following transportation (20). In addition, haematological and serum biochemical changes occur in horses confined during air transport (13). In the present study significantly higher levels in glucose, total protein, albumin, AST and CK levels ( $P < .05$ ) were detected after transportation.

In horses in general CPK and AST are clinically important with muscular diseases because of their high activity in skeletal muscle (21,22,23). Moderate increases in CPK may be associated with moderate exercise or a training program, indeed more severe exercise and competitions may induce increases over 1.000 IU/L. In this study, the CK elevated following transport, and the activity of AST rose in response to transport. The increase in those two serum enzymes indicates minimal muscular insult due to 19 h of transport.

During stressful conditions as exercise or transportation, the hypothalamic-pituitary-adrenal axis activates and may result with an elevated plasma cortisol. Plasma cortisol concentration exhibited variation, with increase during 19h of transport. After the stressor (i.e., transportation) ceased, cortisol dramatically declined. This may be explained with cortisol's relatively short half-life of 1 to 1.5 h (23). The peak in cortisol levels was greater than post-transit cortisol levels evaluated in slaughter horses traveling for up to 30 h (20) but similar to concentrations in moderately exercised horses (20).

In a previous study of 9 h of transportation increased concentrations of cortisol was reported (3). In racehorses, short-term (300 km) transportation or an exercise effected similar serum enzyme and metabolic changes (5). The increases in serum total protein, Na, K and Cl levels after transportation ( $P < 0.05$ ) indicate that the present horses may have had hypertonic dehydration. Although the horses had had water available in their boxes, during a stressful condition such as transport

dehydration may occur. In addition Hct levels exhibited variations with increases after transportation ( $P<0.05$ ) in this study.

WBC counts were deemed as a general indicator of health and the ability of the horses handling stress during transport. The WBC counts showed increases during 24 hours transport in horses (20), however the WBC counts did not exhibit a variation and remained within the reference range for horses during this study. Regarding other relevant hematological indices RBC, Hb and Hct levels exhibited variations with increases after transportation ( $P<0.05$ ) and MCHC exhibited variations with slight decrease after transportation ( $P<0.05$ ).

As a result of horses undergoing 19 h transportation showed patophysiological responses that involve changes in stress indices, hematological and serum biochemical parameters, and dehydration. Changes in PCV (Ht) values may suggest that the horses did not consume sufficient water. The rise in cortisol concentration following transport may have an influence on the immune system. In addition the alterations in glucose concentration may have effect on energy metabolism.

## REFERENCES

1. Austin SM., Foreman JH., Hungerford LL. 1995., Case-control study of risk factors for development of pleuropneumonia in horses. *J Am Vet Med Assoc*, 207, 325-328.
2. Baucus KL., Ralston SL., Nockels CF., McKinnon AO., Squires EL., 1990. Effects of transport on early embryonic death in mares. *J Anim Sci*, 68, 345-351.
3. Baucus KL., Squires EL., Ralston SL., McKinnon AO., Nett TM., 1990. Effect of transportation on the estrus cycle and concentration of hormones in mares. *J Anim Sci* 68, 419-426.
4. Clark DA., Friend TH., Dellmeier G., 1993. The effect of orientation during trailer transport on heart rate, cortisol and balance in horses. *Appl Anim Behav Sci*, 38, 179-189.
5. Codazza D., Maffeo G., Redaelli G., 1974. Serum enzyme changes and hemato-chemical levels in Thoroughbreds after transport and exercise. *J South African Vet Assoc*, 45, 331-334.
6. Crisman MV., Hodgson DR., Bayly WM., Liggitt HD., 1992. Effects of transport on constituents of bronchoalveolar lavage fluid from horses. *Cornell Vet*, 82, 233-246.
7. Fazio E., Ferlazzo A., 2003. Evaluation of Stress During Transport. *Vet Res Commun*, 27, 519-524.
8. Ferlazzo A., Caola G., Omero A., Panzera M., 1984. Modificazioni sieroenzimatiche nel Cavallo sedentario dopo trasporto. *Arch Vet Ital* 35, 27-30.
9. Friend TH., 2000. Dehydration, stress, and water consumption of horses during long-distance commercial transport. *J Anim Sci*, 78, 2568-2580.
10. Friend TH., Martin MT., Householder DD., Bushong DM., 1988. Stress responses of horses during a long period of transport in commercial truck. *J Am Vet Med Assoc*, 212, 838-844.
11. Hodgson DR., 1990. Diseases of the Muscle. In: Smith, BP. (Ed.) *Large Animal Internal Medicine*, Mosby Co., St.Louis. 4th Edition page 1336.
12. Lassourd V., Gayrard V., Laroute M., Alvinerie M., Benard P., Courtot Toutain PL., 1996. Cortisol deposition and production rate in horses during rest and exercise. *J Am Phys*, 271, 25-33.
13. Leadon DP., Frank C., Backhouse W., 1989. A preliminary report on studies on equine transit stress. *J Equ Vet Sci*, 9, 200-202.
14. Leadon DP., Daykin J., Blackhouse W., Frank C., Atock MA., 1990. Environmental, hematological and biochemical changes in equine transit stress. *Proceedings of the Am Assoc Equine Pract*, 36, 465-490.
15. Munro C., Lasley B., 1988. Non-radiometric methods for immunoassay of steroids hormones. In: Albertson, B.D. and H.P. (Ed.) *Non-radiometric Assays: Technology and Application in Polypeptide and Steroid Hormone Detection*. New York. Page 289-329 Alan R Liss, New York.
16. Munro C., Stabenfeldt G., 1984. Development of

- cortisol enzyme immunoassay in plasma. *J Endocrinol*, 101, 41-49.
17. Oikawa M., Kamada M., Toshikawa Y., Yoshikawa T., 1994. Pathology of equine pneumonia associated with transport and isolation of *Streptococcus equi* subsp. *zooepidemicus*. *J Comp Path*, 111, 205-212.
  18. Owen RR., Fullerton J., Barnum DA., 1983. Effects of transportation, surgery, and antibiotic therapy in ponies infected with *Salmonella*. *American J Vet Res*, 44, 46-50.
  19. Stull CL., Rodiek AV., 1995. Effects of post prandial interval and feed type on substrate availability during exercise. *Equine Vet J*, 27, 362-366.
  20. Stull CL., Rodiek AV., 2000. Physiological responses of horses to 24 hours of transportation using a commercial van during summer conditions. *J Anim Sci*, 78, 1458-1466.
  21. Stull CL., Rodiek AV., 2002. Effects of cross-tying horses during 24 h of road transport. *Equine Vet J*, 34, 550-555.
  22. Waran NK., Cuddeford D., 1995. Effects of loading and transport on the heart rate and behavior of horses. *App Anim Behav Sci*, 43, 71-81.
  23. Waran N., Leadon D., Friend T., 2007. The effects of transportation on the welfare of horses. In: Waran N. (Ed.). *The Welfare of Horses*. Kluwer Academic Publishers: Springer, 1, 125-150.
  24. Wessely SJ., Bełkot Z., Bobowiec R., Kosior KU., Wojcik M., 2015. Transport induced inflammatory responses in horses *J Polish Vet Sci*, 18, 407-413.
  25. Nemeč Svete A., Cebulj-Kadunc N., Frangež R., Kruljč P., 2012. Serum cortisol and haematological, biochemical and antioxidant enzyme variables in horse blood sampled in a slaughterhouse lairage, immediately before stunning and during exsanguination. *J Anim Sci*, 6, 1300-1306.
  26. Wernicki A., Urban-Chmiel R., Kankofer M., Mikucki P., Puchalski A., Tokarzewski S., 2006. Evaluation of plasma cortisol and TBARS levels in calves after short-term transportation. *Rev Med Vet*, 157, 30-34.
  27. Andronie I., Parvu M., Andronie V., Ciurea A., 2009. Effects of transportation stress on some physiological indicators in sport horses. *J Zootech Bio*, 42, 379-384.
  28. Onmaz AC., Van den Hoven R., Gunes V., Cinar M., Kucuk O., 2011. Oxidative stress in horses after a 12-hours transport period. *Rev Med Vet*, 162, 213-217.
  29. Tateo A., Padalino B., Boccaccio M., Maggiolino A., Centoducati P., 2012. Transport stress in horses: effects of two different distances. *J Vet Behav Clin App Res*, 7, 33-42.